
AS and A-level Mathematics

Teaching Guidance

AS 7356 and A-level 7357

For teaching from September 2017
For AS and A-level exams from June 2018

Version 1.0, May 2017

Our specification is published on our website (aga.org.uk). We will let centres know in writing about any changes to the specification. We will also publish changes on our website. The definitive version of our specification will always be the one on our website and may differ from printed versions.

You can download a copy of this teaching guidance from our All About Maths website (allaboutmaths.agas.org.uk/). This is where you will find the most up-to-date version, as well as information on version control.

Contents

		Key
General information – disclaimer	5	Pure maths
Subject content	5	Mechanics
A Proof	6	Statistics
B Algebra and functions	8	
C Coordinate geometry in the (x, y) plane	32	
D Sequences and series	39	
E Trigonometry	50	
F Exponentials and logarithms	69	
G Differentiation	80	
H Integration	96	
I Numerical methods	112	
J Vectors	119	
For sections K to O students must demonstrate the ability to use a calculator to compute summary statistics and access probabilities from standard statistical distributions.		
K Statistical sampling	127	
L Data presentation and interpretation	131	
M Probability	143	
N Statistical distributions	151	
O Statistical hypothesis testing	158	
P Quantities and units in mechanics	163	
Q Kinematics	165	
R Forces and Newton's laws	179	
S Moments	194	
1 Appendix A Mathematical notation for AS and A-level qualifications in Mathematics and Further Mathematics	196	
2 Appendix B Mathematical formulae and identities	205	

General information – disclaimer

This AS and A-level Mathematics teaching guidance will help you plan your teaching by further explaining how we have interpreted content of the specification and providing examples of how the content of the specification may be assessed. The teaching guidance notes do not always cover the whole content statement.

The examples included in this guidance have been chosen to illustrate the level at which this content will be assessed. The wording and format used in this guidance do not always represent how questions would appear in a question paper. Not all questions in this guidance have been through the same rigorous checking process as the ones used in our question papers.

Several questions have been taken from legacy specifications and therefore represent higher levels of AO1 than will be found in a suite of exam papers for this A-level Mathematics specification.

This guidance is not, in any way, intended to restrict what can be assessed in the question papers based on the specification. Questions will be set in a variety of formats including both familiar and unfamiliar contexts.

All knowledge from the GCSE Mathematics specification is assumed.

Subject content

The subject content for AS and A-level Mathematics is set out by the Department for Education (DfE) and is common across all exam boards.

This document is designed to illustrate the detail within the content defined by the DfE.

Content in **bold type** is contained within the AS Mathematics qualification as well as the A-level Mathematics qualification. Content in standard type is contained only within the A-level Mathematics qualification.

A

Proof

A1

Understand and use the structure of mathematical proof, proceeding from given assumptions through a series of logical steps to a conclusion; use methods of proof, including proof by deduction, proof by exhaustion.

Disproof by counter example.

Proof by contradiction (including proof of the irrationality of $\sqrt{2}$ and the infinity of primes, and application to unfamiliar proofs).

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- set out a clear proof with the correct use of symbols, such as $=$, \Rightarrow , \Leftarrow , \Leftrightarrow , \equiv , \therefore , \because
- understand that many examples can be useful in looking for structure, but they do not constitute a proof.

Note: at A-level 25% (20% at AS) of the assessment material must come from Assessment Objective 2 (reason, interpret and communicate mathematically). A focus on clear reasoning, mathematical argument and proof using precise mathematical language and notation should underpin the teaching of this specification. Students should become familiar with the mathematical notation found in Appendix A of the specification.

Examples

- 1 Prove that 113 is a prime number.
- 2 Prove that for any positive whole number, n , the value of $n^3 - n$ is always a multiple of 3.
- 3 “For any positive whole number, n , the value of $2n^2 + 11$ is a prime number.”
Find a value for n that disproves this statement.

Only assessed at A-level

Examples

- 1 Assuming $\sqrt{2}$ is a rational number we can write $\sqrt{2} = \frac{a}{b}$, where a and b are positive whole numbers with no common factors.
- (a) Show that a must be even.
 - (b) Show that b must be even.
 - (c) Using parts (a) and (b), explain why there is a contradiction and state what conclusion can be made about $\sqrt{2}$ as a result.

B

Algebra and functions

B1

Understand and use the laws of indices for all rational exponents.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand and use the following laws:

$$x^a \times x^b = x^{a+b}$$

$$x^a \div x^b = x^{a-b}$$

$$(x^a)^b = x^{ab}$$

$$x^{-a} = \frac{1}{x^a}$$

$$x^{\frac{a}{b}} = \sqrt[b]{x^a} = (\sqrt[b]{x})^a$$

- apply these laws when solving problems in other contexts, for example simplification of expressions before integrating/differentiating, solving equations or transforming graphs.

Examples

- 1 (a) Write down the values of p , q and r , given that:

(i) $64 = 8^p$

(ii) $\frac{1}{64} = 8^q$

(iii) $\sqrt{8} = 8^r$

- (b) Find the value of x for which $\frac{8^x}{\sqrt{8}} = \frac{1}{64}$

- 2 Find $\int \left(x + 1 + \frac{4}{x^2} \right) dx$

3 A curve has equation $y = \frac{1}{x^2} + 4x$

Find $\frac{dy}{dx}$

B2

Use and manipulate surds, including rationalising the denominator.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- demonstrate they understand how to manipulate surds and rationalise denominators.
- show a result.
- answer algebraic questions.

Note: many calculators will perform simplifications but students should understand that the command words 'show that' require detailed mathematical reasoning using precise notation.

Examples

1 Show that $\frac{\sqrt{75} - \sqrt{27}}{\sqrt{3}}$ is an integer and find its value.

2 Rationalise the denominator of the fraction $\frac{3 + \sqrt{a}}{2 - \sqrt{a}}$, where a is a positive integer.

3 A rectangle has length $(9 + 5\sqrt{3})$ cm and area $(15 + 7\sqrt{3})$ cm²

Show that the width of the rectangle is $(m + n\sqrt{3})$, where m and n are integers.

4 (a) Expand $(\sqrt{x} - 1)^2$

(b) Hence find $\int (\sqrt{x} - 1)^2 dx$

5 (a) Write $\sqrt{x^5}$ in the form x^k where k is a fraction.

(b) Find $\int (7\sqrt{x^5} - 4) dx$

B3

Work with quadratic functions and their graphs; the discriminant of a quadratic function, including the conditions for real and repeated roots; completing the square; solution of quadratic equations including solving quadratic equations in a function of the unknown.

Assessed at AS and A-level



Teaching guidance

Students should:

- be able to sketch graphs of quadratics, ie of $y = ax^2 + bx + c$
- be able to identify features of the graph such as points where the graph crosses the axes, lines of symmetry or the vertex of the graph.
- know and use the following:



$$b^2 - 4ac > 0$$

Distinct real roots





$$b^2 - 4ac = 0$$

Equal roots



$$b^2 - 4ac < 0$$

No real roots



Note: a quadratic described as having real roots will be such that $b^2 - 4ac \geq 0$

- be able to complete the square and use the resulting expression to make deductions, such as the maximum/minimum value of a quadratic or the number of roots.

Note: unless specified in a question, any correct method for solving a quadratic will be acceptable. Students should become familiar with using calculators to solve quadratic equations. Use of appropriate technology should be encouraged throughout the teaching of this specification. It is a requirement from the DfE that 'the use of technology...must permeate the study of AS and A-level Mathematics'.

Teaching guidance continued

- be able to solve quadratic equations in a function of the unknown, where the function may be, for example, trigonometric or exponential.

Note: quadratic equations may arise from problems set in a variety of contexts taken from mechanics and statistics. The context may determine the level of the question (AS or A-level).

Examples

- Express $x^2 + 8x + 19$ in the form $(x + p)^2 + q$, where p and q are integers.
 - Hence, or otherwise, show that the equation $x^2 + 8x + 19 = 0$ has no real solutions.
 - Sketch the graph of $y = x^2 + 8x + 19$, stating the coordinates of the minimum point and the point where the graph crosses the y -axis.
- Find the values of k for which the equation $x^2 - 2(k + 1)x + 2k^2 - 7 = 0$ has equal roots.
- Factorise $9 - 8x - x^2$
 - Show that

$$25 - (x + 4)^2$$

can be written as

$$9 - 8x - x^2$$
 - A curve has equation $y = 9 - 8x - x^2$
 - Write down the equation of its line of symmetry.
 - Find the coordinates of its vertex.
 - Sketch the curve, indicating the values of the intercepts on the x -axis and the y -axis.

- 4 (a) Using the substitution $Y = 3^x$, show that the equation

$$9^x - 3^{x+1} + 2 = 0$$

can be written as

$$(Y - 1)(Y - 2) = 0$$

- (b) Hence show that the equation

$$9^x - 3^{x+1} + 2 = 0$$

has a solution

$$x = 0$$

and, by using logarithms, find the other solution, giving your answer to four decimal places.

- 5 Given that

$$\frac{3 + \sin^2 \theta}{\cos \theta - 2} = 3 \cos \theta$$

show that

$$\cos \theta = -\frac{1}{2}$$

- 6 Solve the equation

$$3^{2x} - 3^{x+1} - 4 = 0$$

giving your answer in an exact form.

B4

Solve simultaneous equations in two variables by elimination and by substitution, including one linear and one quadratic equation.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand the relationship between the algebraic solution of simultaneous equations and the points of intersection on the corresponding graphs.
- in the case of one linear and one quadratic equation, recognise the geometrical significance of the discriminant of the resulting quadratic.

Note: simultaneous equations could arise from problems set on a variety of topics including mechanics and statistics.

Examples

1 The straight line L has equation

$$y = 3x - 1$$

The curve C has equation

$$y = (x + 3)(x - 1)$$

- Sketch on the same axes the line L and the curve C , showing the values of the intercepts on the x -axis and y -axis.
- Show that the x -coordinates of the points of intersection L and C satisfy the equation $x^2 - x - 2 = 0$
- Hence find the coordinates of the points of intersection of L and C .

2 The curve C has equation

$$y = k(x^2 + 3)$$

where k is a constant.

The line L has equation

$$y = 2x + 2$$

Show that the x -coordinates of any points of intersection of the curve C with the line L satisfy the equation

$$kx^2 - 2x + 3k - 2 = 0$$

- 3 A circle with centre C has equation

$$x^2 + y^2 - 10y + 20 = 0$$

A line has equation

$$y = 2x$$

- (a) Show that the x -coordinate of any point of intersection of the line and the circle satisfies the equation $x^2 - 4x + 4 = 0$
- (b) Hence, show that the line is a tangent to the circle and find the coordinates of the point of contact P .
- 4 The first term of an arithmetic series is 1. The common difference of the series is 6.
- (a) Find the 10th term of the series.
- (b) The sum of the first n terms of the series is 7400.
- (i) Show that $3n^2 - 2n - 7400 = 0$
- (ii) Find the value of n .

B5

Solve linear and quadratic inequalities in a single variable and interpret such inequalities graphically, including inequalities with brackets and fractions.

Express solutions through correct use of 'and' and 'or', or through set notation.

Represent linear and quadratic inequalities such as $y > x + 1$ and $ax^2 + bx + c$ graphically.

Assessed at AS and A-level

Teaching guidance

Students should be able to:


- give the range of values which satisfy more than one inequality.
- illustrate regions on sketched graphs, defined by inequalities.
- define algebraically inequalities that are given graphically.

Notes: Dotted/dashed lines or curves will be used to indicate strict inequalities.

Overarching theme 1.3 is of particular relevance here. Students are required to demonstrate an understanding of and use the notation (language and symbols) associated with set theory (as set out in Appendix A of the specification). Students are required to apply this notation to the solutions of inequalities.

Examples

- 1 Find the possible values of k which satisfy the inequality $3k^2 - 2k - 1 < 0$
- 2 The diagram shows the graphs of $y = x + 5$ and $y = x^2 - 4x + 5$.



State which pair of inequalities defines the shaded region.

Circle your answer.

$$\begin{array}{c} y < x + 5 \\ \text{and} \\ y < x^2 - 4x + 5 \end{array}$$

$$\begin{array}{c} y \leq x + 5 \\ \text{or} \\ y > x^2 - 4x + 5 \end{array}$$

$$\begin{array}{c} y \leq x + 5 \\ \text{and} \\ y \geq x^2 - 4x + 5 \end{array}$$

$$\begin{array}{c} y \geq x + 5 \\ \text{or} \\ y < x^2 - 4x + 5 \end{array}$$

- 3 Find the values for x which satisfy both inequalities $x^2 + 2x > 8$ and $3(2x + 1) \leq 15$.

B6

Manipulate polynomials algebraically, including expanding brackets and collecting like terms, factorisation and simple algebraic division; use of the factor theorem.

Simplify rational expressions including by factorising and cancelling, and algebraic division (by linear expressions only).

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- manipulate polynomials, which may be embedded in questions focused on other topics.
- understand factorisation and division applied to a quadratic or a cubic polynomial divided by a linear term of the form $(x + a)$ where a is an integer.

Notes: Any correct method will be accepted, eg by inspection, by equating coefficients or by formal division.

The greatest level of difficulty is exemplified by $x^3 - 5x^2 + 7x - 3$, ie a cubic always with a factor $(x + a)$, where a is a small integer and including the cases of three distinct linear factors, repeated linear factors or a quadratic factor which cannot be factorised in the real numbers.

Examples

1 Find $\int (2x+1)(x^2 - x + 2)dx$

2 The polynomial $p(x)$ is given by $p(x) = x^3 + 7x^2 + 7x - 15$

- Use the Factor Theorem to show that $x + 3$ is a factor of $p(x)$
- Express $p(x)$ as the product of three linear factors.

3 The polynomial $p(x)$ is given by $p(x) = x^3 + x - 10$

- Use the Factor Theorem to show that $x - 2$ is a factor of $p(x)$
- Express $p(x)$ in the form $(x - 2)(x^2 + ax + b)$, where a and b are constants.

Only assessed at A-level

Teaching guidance

Students should be able to:

- understand the Factor Theorem where the divisor is of the form $(ax + b)$.
- simplify rational expressions.
- carry out algebraic division where the divisor is of the form $(ax + b)$.

Note: any correct method will be accepted, eg by inspection, by equating coefficients or by formal division.

Examples

1 Express $\frac{3x^3 + 8x^2 - 3x - 5}{3x - 1}$ in the form $ax^2 + bx + \frac{c}{3x - 1}$, where a , b and c are integers.

2 The polynomial $f(x)$ is defined by $f(x) = 4x^3 - 7x - 3$

(a) Find $f(-1)$

(b) Use the Factor Theorem to show that $2x + 1$ is a factor of $f(x)$

(c) Simplify the algebraic fraction $\frac{4x^3 - 7x - 3}{2x^2 + 3x + 1}$

B7

Understand and use graphs of functions; sketch curves defined by simple equations including polynomials, the modulus of a linear function, $y = \frac{a}{x}$ and $y = \frac{a}{x^2}$ (including their vertical and horizontal asymptotes); interpret algebraic solution of equations graphically; use intersection points of graphs to solve equations.

Understand and use proportional relationships and their graphs.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand, use and sketch straight-line graphs (including vertical and horizontal).
- understand and use polynomials up to cubic (including sketching curves).
- understand and use cubic polynomials with at least one linear factor.
- distinguish between the various possibilities for graphs of cubic polynomials indicating where graphs meet coordinate axes.
- understand and use graphs of the functions $y = \frac{a}{x}$ and $y = \frac{a}{x^2}$ as well as simple transformations of these graphs (including sketching curves).
- use the following:


Proportionality (\propto)	Equation
y is proportional to x	$y = kx$
y is proportional to x^n	$y = kx^n$
y is inversely proportional to x^n	$y = \frac{k}{x^n}$

Examples

- 1 (a) Sketch the graph of $y = \frac{1}{x}$
- (b) State the equations of the asymptotes.
- (c) Sketch the graph of $y = \frac{1}{x+2}$ clearly labelling the vertical asymptote.
- (d) State the transformation which maps the graph of $y = \frac{1}{x+2}$ on to $y = \frac{1}{x}$

It should be noted that these sketches can be obtained from many calculators so relatively little credit will be given for the sketch and no credit will be given for a poor sketch. Students must also understand the importance of labelling diagrams and sketches.

- 2 The graphs of $y = x^3 - ax^2 + b$ and $y = cx + d$ are shown on the diagram.



One of the points of intersection of the two graphs is $(3, 4)$.

Find the values of a , b , c and d .

Only assessed at A-level

Teaching guidance

Students should:

- know and be able to use:


$$|x| = \begin{cases} x & \text{for } x \geq 0 \\ -x & \text{for } x < 0 \end{cases}$$

- understand and be able to use the graph of $y = |x|$ and combinations of simple transformations of this graph, points of intersection and solutions of equations and inequalities.

Examples

- Sketch the graph of $y = |2x|$
 - On a separate diagram, sketch the graph of $y = 4 - |2x|$ including the coordinates of the points where the graph crosses the coordinate axes.
 - Solve $4 - |2x| = x$
 - Hence, or otherwise, solve the inequality $4 - |2x| > x$
- Sketch the graph of $y = |8 - 2x|$
 - Solve the inequality $|8 - 2x| > 4$

- 3 The diagram below shows the graphs of $y = |2x - 3|$ and $y = |x|$



- (a) Find the coordinates of the points of intersection of the graphs of $y = |2x - 3|$ and $y = |x|$
- (b) Hence, or otherwise, solve the inequality $|2x - 3| \geq |x|$

B8

Understand and use composite functions, inverse functions, and their graphs.

Only assessed at A-level

Teaching guidance

Students should:

- be able to define a function as one-to-one or many-to-one mapping, including the range and domain (co-domain not required).
- understand that the domain and range of a function are sets.
- understand and be able to use correct language and notation to describe functions accurately.
- know the conditions for the existence of the inverse and the relationship between a function's range and domain, and that of the inverse function.
- understand that f^{-1} is the inverse of $f \Leftrightarrow f^{-1}f(x) = x$ for all x
- recognise and be able to use notation such as:
 - $f : x \mapsto y$
 - $f(x) = x^2$
 - f^{-1} to indicate inverse.
- understand that the graph of an inverse function can be found by reflecting in the line $y = x$
- understand the composition of functions:
 $fg(x)$ is f applied to the result of $g(x)$
 and know that the range of g must be a subset of the domain of f .

Note: questions relating to this section of content will be set in the context of overarching theme 1.4

Examples

- 1 The functions f and g are defined with their respective domains by

$$f(x) = e^{2x} - 3, \text{ for all real values of } x.$$

$$g(x) = \frac{1}{3x+4}, \text{ for all real values of } x, x \neq -\frac{4}{3}$$

- (a) Find the range of f .


- (b) The inverse of f is f^{-1}

$$\text{Solve the equation } f^{-1}(x) = 0$$

- (c) (i) Find an expression for $gf(x)$
- (ii) Solve the equation $gf(x) = 1$, giving your answer in exact form.
- (iii) For what value of x is $gf(x)$ undefined?

- 2 The curve with equation $y = f(x)$, where $f(x) = \ln(2x - 3)$, is sketched below.

The domain of f is $\{x \in \mathbb{R} : x > \frac{3}{2}\}$



- (a) The inverse of f is f^{-1}
- Find $f^{-1}(x)$
 - State the range of f^{-1}
 - Sketch the curve with equation $y = f^{-1}(x)$, indicating the value of the y -coordinate of the point where the curve intersects the y -axis.
- (b) The function g is defined by $g(x) = e^{2x} - 4$, for all real values of x
- Find $gf(x)$, giving your answer in the form $(ax - b)^2 - c$, where a , b and c are integers
 - Write down an expression for $fg(x)$, and hence find the exact solution of the equation $fg(x) = \ln 5$

B9

Understand the effect of simple transformations on the graph of $y = f(x)$ including sketching associated graphs: $y = af(x)$, $y = f(x) + a$, $y = f(x + a)$ and $y = f(ax)$ and combinations of these transformations.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- describe the following:

$y = af(x)$	Stretch in the y -direction scale factor a
$y = f(x) + a$	Translation by the vector $\begin{pmatrix} 0 \\ a \end{pmatrix}$
$y = f(x + a)$	Translation by the vector $\begin{pmatrix} -a \\ 0 \end{pmatrix}$
$y = f(ax)$	Stretch in the x -direction scale factor $\frac{1}{a}$

- understand that applying different transformations may result in the same function.

Examples

1 Describe a single geometrical transformation that maps the graph of $y = 3^x$

- (a) onto the graph of $y = 3^{2x}$
- (b) onto the graph of $y = 3^{x+1}$

2 The graph of $y = x^2 - 6x + 9$ is translated by the vector $\begin{pmatrix} -3 \\ 0 \end{pmatrix}$

Find the equation of the translated graph, giving your answer in its simplest form.

3 The graph of $y = 2^x$ is mapped by a single transformation onto the graph of $y = 2^{x-2}$

(a) Fully describe the single transformation as a translation.

(b) Fully describe the single transformation as a stretch.

Only assessed at A-level

Teaching guidance

Students should be able to:

- apply two or more transformations to a function or describe a combination of two or more transformations that result in a given function.
- understand that applying transformations in a different order may result in two different functions.

Example

- 1 Describe a sequence of two geometrical transformations that maps the graph of $y = \cos^{-1} x$ onto the graph of $y = 2\cos^{-1}(x-1)$

B10

Decompose rational functions into partial fractions (denominators not more complicated than squared linear terms and with no more than three terms, numerators constant or linear).

Only assessed at A-level

Teaching guidance

Students should be able to:

- use the following forms:

$$\frac{px+q}{(ax+b)(cx+d)(ex+f)} \equiv \frac{A}{(ax+b)} + \frac{B}{(cx+d)} + \frac{C}{(ex+f)}$$

$$\frac{px+q}{(ax+b)(cx+d)^2} \equiv \frac{A}{(ax+b)} + \frac{B}{(cx+d)} + \frac{C}{(cx+d)^2}$$

- understand that the fractions may need to be simplified before partial fractions are found.
- understand that partial fractions may be required for integration or in a binomial approximation.

Examples

- 1 (a) Express $\frac{2x+3}{4x^2-1}$ in the form $\frac{A}{2x-1} + \frac{B}{2x+1}$ where A and B are integers.
- (b) Express $\frac{12x^3-7x-6}{4x^2-1}$ in the form $Cx + \frac{D(2x+3)}{4x^2-1}$ where C and D are integers.
- (c) Evaluate $\int_1^2 \frac{12x^3-7x-6}{4x^2-1} dx$ giving your answer in the form $p + \ln q$, where p and q are rational numbers.

2 (a) Express $\frac{1}{(3-2x)(1-x)^2}$ in the form $\frac{A}{3-2x} + \frac{B}{1-x} + \frac{C}{(1-x)^2}$

(b) Solve the differential equation

$$\frac{dy}{dx} = \frac{2\sqrt{y}}{(3-2x)(1-x)^2}$$

where $y = 0$ when $x = 0$, expressing your answer in the form

$$y^p = q \ln[f(x)] + \frac{x}{1-x}$$

where p and q are constants.

3 It is given that $f(x) = \frac{7x-1}{(1+3x)(3-x)}$

(a) Express $f(x)$ in the form $\frac{A}{3-x} + \frac{B}{1+3x}$ where A and B are integers.

(b) (i) Find the first three terms of the binomial expansion of $f(x)$ in the form $a + bx + cx^2$, where a , b and c are rational numbers.

(ii) State why the binomial expansion cannot be expected to give a good approximation of $f(x)$ at $x = 0.4$

B11

Use of functions in modelling, including consideration of limitations and refinements of the models.

Only assessed at A-level

Teaching guidance

Students should be able to

- suggest how a model could be improved.
- give suggestions as to when a particular model might break down or why it is only appropriate over a particular range of values.

Note: functions may be used in the formulation of a differential equation or arise from the solution of a differential equation.

Examples

- 1 The total number of views, V , of a viral video clip that is released on the internet is given by the formula

$$V = 150 \times 2^d$$

where d is the number of days after it has been released.

- (a) How many days does it take for the video to reach 1 million hits?
- (b) Explain why this model will eventually break down.

- 2 A zoologist is studying a population of 100 rodents introduced on to a small island. In order to model the size of the population, she assumes that the rate of increase of the number of rodents, $\frac{dN}{dt}$, at time t , will be proportional to the size of the population, N .

This leads the zoologist to the model:

$$N = Ae^{kt}$$

- (a) State the value of A .
- (b) Describe what happens to the population of rodents as time increases.
- (c) State one criticism of this model and explain how it could be improved.

Note: A-level students should be encouraged to apply their mathematical thinking to problem solving and modelling from the first day of teaching.

C

Coordinate geometry in the (x, y) plane

C1

Understand and use the equation of a straight line, including the forms $y - y_1 = m(x - x_1)$ and $ax + by + c = 0$; gradient conditions for two straight lines to be parallel or perpendicular.

Use straight line models in a variety of contexts.

Assessed at AS and A-level

Teaching guidance

Students should:


- be able to solve problems using gradients, midpoints and the distance between two points, including the form $y = mx + c$ and the forms $y = a$ and $x = a$ for horizontal and vertical lines.
- know that the product of the gradients of two perpendicular lines is -1 .

Note: in questions where the equation of a line is to be found, any correct form will be acceptable, unless specified in the question. However, trivial simplifications left undone in equations are likely to

be penalised, eg $y - -2 = \frac{2}{4}(x - 1)$ should be simplified to $y + 2 = \frac{1}{2}(x - 1)$

Examples

- 1 The trapezium $ABCD$ is shown below.



The line AB has equation $2x + 3y = 14$ and DC is parallel to AB .

- (a) Find the gradient of AB .
- (b) The point D has coordinates $(3, 7)$
 - (i) Find an equation of the line DC .
 - (ii) The angle BAD is a right angle.
Find an equation of the line AD , giving your answer in the form $mx + ny + p = 0$, where m , n and p are integers.
- (c) The line BC has equation $5y - x = 6$. Find the coordinates of B .

- 2 The point A has coordinates $(-1, 2)$ and the point B has coordinates $(3, -5)$

- (a)
 - (i) Find the gradient of AB .
 - (ii) Hence find an equation of the line AB , giving your answer in the form $px + qy = r$, where p , q and r are integers.
- (b) The midpoint of AB is M .
 - (i) Find the coordinates of M .
 - (ii) Find an equation of the line which passes through M and which is perpendicular to AB .
- (c) The point C has coordinates $(k, 2k + 3)$.

Given that the distance from A to C is $\sqrt{13}$ find the two possible values of the constant k .

C2

Understand and use the coordinate geometry of the circle including using the equation of a circle in the form $(x - a)^2 + (y - b)^2 = r^2$; completing the square to find the radius of a circle; use the following properties:

- the angle in a semicircle is a right angle
- the perpendicular from the centre to a chord bisects the chord
- the radius of a circle at a given point on its circumference is perpendicular to the tangent to the circle at that point.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- find the equation of a tangent or normal at a point.
- find relevant gradients using the coordinates of appropriate points.

Note: implicit differentiation will not be required at AS.

Examples

1 A circle with centre C has equation $x^2 + y^2 - 10x + 12y + 41 = 0$

The point $A(3, -2)$ lies on the circle.

(a) Express the equation of the circle in the form

$$(x - a)^2 + (y - b)^2 = k$$

(b) (i) Write down the coordinates of C .

(ii) Show that the circle has radius $n\sqrt{5}$, where n is an integer.

(c) Find the equation of the tangent to the circle at point A , giving your answer in the form $x + py = q$, where p and q are integers.

(d) The point B lies on the tangent to the circle at A and the length of BC is 6.
Find the length of AB .

2 The points $P(4, 3)$, $Q(6, 7)$ and $R(12, 4)$ lie on a circle, C .

- (a) Show that PQ and QR are perpendicular
- (b) Find the length of PR , giving your answer as a surd.
- (c) Find the equation of the circle C .

3 A circle has equation $x^2 + y^2 - 4x - 14 = 0$

- (a)
 - (i) Find the coordinates of the centre of the circle.
 - (ii) Find the radius of the circle in the form $p\sqrt{2}$, where p is an integer.
- (b) A chord of the circle has length 8. Find the perpendicular distance from the centre of the circle to this chord.
- (c) A line has equation $y = 2k - x$ where k is a constant.

- (i) Show that the x -coordinate of any point of intersection of the line and the circle satisfies the equation

$$x^2 - 2(k + 1)x + 2k^2 - 7 = 0$$

- (ii) Find the values of k for which the equation

$$x^2 - 2(k + 1)x + 2k^2 - 7 = 0$$

has equal roots.

- (iii) Describe the geometrical relationship between the line and the circle when k takes either value found in part (c)(ii).

C3

Understand and use the parametric equations of curves and conversion between Cartesian and parametric forms.

Only assessed at A-level

Teaching guidance

Students should:

- understand that it is not always expected that the resulting Cartesian equation will be in explicit form.
Note: any correct form will be acceptable, unless stated in the question.
- understand that parametric equations using trigonometric terms, often require the use of appropriate identities.
- be able to answer questions requiring gradient of, or tangent to a curve given in parametric form.

Examples

- 1 A curve is defined by the parametric equations

$$x = 3 - 4t \quad y = 1 + \frac{2}{t}$$

Verify that the Cartesian equation of the curve can be written as

$$(x - 3)(y - 1) + 8 = 0$$

- 2 A curve is given by the parametric equations


$$x = \cos \theta \quad y = \sin 2\theta$$

Show that the Cartesian equation of the curve can be written as

$$y^2 = kx^2(1 - x^2)$$

where k is an integer.

- 3 The curve with parametric equations $y = 1 - \cos \theta$ and $x = \sin \theta, 0 \leq \theta \leq 2\pi$ is shown below.



Find a Cartesian equation of the curve.

C4

Use parametric equations in modelling in a variety of contexts.

Only assessed at A-level

Teaching guidance


Students should be able to use parametric equations to describe the motion of a particle in the (x, y) plane, for example $x = 4t$, $y = 3t - 4.9t^2$, using the acceleration due to gravity as 9.8 m/s^2 for a particle subject

only to the force of its own weight, projected from the origin at time $t = 0$ with velocity $\begin{pmatrix} 4 \\ 3 \end{pmatrix}$

Examples

- 1 A ball is thrown from an origin in a strong wind so that its horizontal and vertical position at time t is given by $x = 25t - 10t^2$ and $y = 15t - 10t^2$

The trajectory of the ball is shown in the diagram.



- (a) Find the time it takes for the ball to hit the ground.
- (b) Find $\frac{dy}{dx}$ in terms of t
- (c) Use your answers to parts (a) and (b) to find the angle at which the ball hits the ground.

Note: A-level students should be encouraged to apply their mathematical thinking to problem solving and modelling from the first day of teaching.

D

Sequences and series

D1

Understand and use the binomial expansion of $(a + bx)^n$ for positive integer n ; the notations $n!$ and nCr ; link to binomial probabilities.

Extend to any rational n , including its use for approximation; be aware that the expansion is valid for $\left|\frac{bx}{a}\right| < 1$ (proof not required).

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- answer questions requiring the full expansion of expressions of the form $(a + bx)^n$, where n is a small positive integer.
- find the coefficients of particular powers of x (complete expansion not required).

Notes: The notation $\binom{n}{r}$ will be acceptable but not required.

The x in $(a + bx)^n$ may be a simple function of x , eg $\left(2 - \frac{1}{x}\right)^4$

Examples

1 (a) Using the binomial expansion, or otherwise:

(i) express $(1 + x)^3$ in ascending powers of x .

(ii) express $(1 + x)^4$ in ascending powers of x .

(b) Hence, or otherwise

(i) express $(1 + 4x)^3$ in ascending powers of x .

(ii) express $(1 + 3x)^4$ in ascending powers of x .

(c) Show that the expansion of

$$(1 + 3x)^4 - (1 + 4x)^3$$

can be written in the form

$$px^2 + qx^3 + rx^4$$

where p , q and r are integers.

2 (a) (i) Using the binomial expansion, or otherwise, express $(2 + y)^3$ in the form

$$a + by + cy^2 + y^3$$

where a , b and c are integers.

(ii) Hence show that

$$(2 + x^{-2})^3 + (2 - x^{-2})^3$$

can be expressed in the form

$$p + qx^{-4}$$

where p and q are integers.

(b) Hence find the value of $\int_1^2 \left[(2 + x^{-2})^3 + (2 - x^{-2})^3 \right] dx$

Only assessed at A-level

Teaching guidance

Students should be familiar with and be able to use the formula:

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{R})$$

where: $\binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!}$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1.2}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{1.2\dots r}x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

Examples

- 1 (a) (i) Find the binomial expansion of $(1-x)^{-1}$ up to the term in x^3
- (ii) Hence, or otherwise, obtain the binomial expansion of $\frac{1}{1+3x}$ up to the term in x^3
- (b) Express $\frac{1+4x}{(1+x)(1+3x)}$ in partial fractions.
- (c) (i) Find the binomial expansion of $\frac{1+4x}{(1+x)(1+3x)}$ up to the term in x^3
- (ii) Find the range of values of x for which the binomial expansion of $\frac{1+4x}{(1+x)(1+3x)}$ is valid.
- 2 (a) Find the binomial expansion of $(1+6x)^{\frac{2}{3}}$ up to and including the term in x^2
- (b) Find the binomial expansion of $(8+6x)^{\frac{2}{3}}$ up to and including the term in x^2
- (c) Use your answer from part (b) to find an estimate for $\sqrt[3]{100}$ in the form $\frac{a}{b}$, where a and b are integers.

D2

Work with sequences including those given by a formula for the n th term and those generated by a simple relation of the form $x_{n+1} = f(x_n)$; increasing sequences; decreasing sequences; periodic sequences.

Only assessed at A-level

Teaching guidance

Students should:

- understand and be able to use notation such as u_n
- understand that an increasing sequence will be one where $u_{n+1} > u_n$ for all n

eg

$$u_{n+1} = 2u_n, u_1 = 5$$

$$u_n = \frac{n}{n+1}$$

- understand that a decreasing sequence will be one where $u_{n+1} < u_n$, for all n

eg

$$u_{n+1} = 0.5u_n, u_1 = 5$$

$$u_n = 2^{-n}$$

- know that a periodic sequence repeats over a fixed interval, ie $u_{n+a} = u_n$

eg

1, 2, 3, 4, 1, 2, 3, 4, ...

$$u_n = \sin\left(\frac{\pi n}{2}\right)$$

- be able to find a limit, L , as $n \rightarrow \infty$ by putting $L = f(L)$

Examples

- 1 The n th term of a sequence is u_n

The sequence is defined by

$$u_{n+1} = pu_n + q$$

where p and q are constants.

The first three terms of the sequence are given by:

$$u_1 = 200$$

$$u_2 = 150$$

$$u_3 = 120$$

- (a) Show that $p = 0.6$ and find the value of q .
- (b) Find the value of u_4 .
- (c) The limit of u_n as n tends to infinity is L . Write down an equation for L and hence find the value of L .
- 2 The n th term of a sequence is defined by

$$u_n = \frac{n}{n+1}$$

- (a) Write down the value of u_1 and u_2 .
- (b) Prove that u_n is an increasing sequence.
- 3 When the fraction $\frac{1250}{999}$ is written as a decimal its digits form a periodic sequence.

What digit is in the 1000th decimal place?

D3

Understand and use sigma notation for sums of series.

Only assessed at A-level

Teaching guidance

Students should be able to:

- use \sum to indicate the sum of a series, eg

$$\sum_{r=1}^5 2r+1 = 3+5+7+9+11$$

$$u_n = 2^{-n}, \sum_{n=0}^{\infty} u_n = 1 + \frac{1}{2} + \frac{1}{4} + \dots = 2$$

- understand and use the notation S_n for the sum of a series.

Note: it should be expected that questions set may require exact forms to avoid trivial calculator solutions.

Examples

- 1 Find the value of

$$\sum_{r=1}^4 \ln(2^r)$$

giving your answer in the form $p \ln 2$.

- 2 Find the exact value of

$$\sum_{r=1}^{\infty} \frac{\sqrt{2}}{2^r}$$

Note: students should be encouraged to read questions carefully and understand that where specified responses are requested alternative forms will not be awarded full credit.

D4

Understand and work with arithmetic sequences and series, including the formulae for n th term and the sum to n terms.

Only assessed at A-level

Teaching guidance

Students should know the formula for the n th term and be able to use the formulae for the sum of the first n terms:

$u_n = a + (n-1)d$	n th term of the sequence
$S_n = \frac{n}{2}(a+l)$	Sum of first n terms using first and last term
$S_n = \frac{n}{2}(2a + (n-1)d)$	Sum of first n terms using first term and common difference

where a is the first term, d is the common difference and l is the last term of the sequence.

Examples

1 The first term of an arithmetic series is 1. The common difference of the series is 6.

- (a) Find the 10th term of the series.
- (b) The sum of the first n terms of the series is 7400.
 - (i) Show that $3n^2 - 2n - 7400 = 0$.
 - (ii) Find the value of n .

2 The 25th term of an arithmetic series is 38.

The sum of the first 40 terms of the series is 1250.

- (a) Show that the common difference of this series is 1.5.
- (b) Find the number of terms in the series which are less than 100.

3 The arithmetic series

$$51 + 58 + 65 + 72 + \dots + 1444$$

has 200 terms.

- (a) Write down the common difference of the series.
- (b) Find the 101st term of the series.
- (c) Find the sum of the last 100 terms of the series.

Note: where one part of a question relies on an answer derived from a previous part ‘follow through’ will be applied in examination assessment, provided clear and correct mathematical method has been applied. Students should be encouraged to explain and justify key steps in their working even where problems are routine in nature.

D5

Understand and work with geometric sequences and series including the formulae for the n th terms and the sum of a finite geometric series; the sum to infinity of a convergent geometric series, including the use of $|r| < 1$; modulus notation.

Only assessed at A-level

Teaching guidance

Students should:

- know the formula for the n th term and be able to use the sum formulae:

$u_n = ar^{n-1}$	n th term of the sequence
$S_n = \frac{a(1-r^n)}{1-r} \left(= \frac{a(r^n-1)}{r-1} \right)$	Sum of first n terms
$S_\infty = \frac{a}{1-r}, r < 1$	Sum to infinity

where a is the first term and r is the common ratio.

- understand the condition for when a geometric series is convergent.

Examples

- 1 A geometric series has first term 80 and common ratio $\frac{1}{2}$.
 - (a) Find the 3rd term of the series.
 - (b) Find the sum to infinity of the series.
 - (c) Find the sum of the first 12 terms of the series, giving your answer to two decimal places.
- 2 The first three terms of a geometric sequence are x , $x + 6$ and $x + 9$.
Find the common ratio of this sequence, giving your answer as a fraction in its simplest form.

-
- 3 An infinite geometric series has common ratio r . The sum to infinity of the series is five times the first term of the series.
- (a) Show that $r = 0.8$.
- (b) Given that the first term of the series is 20, find the least value of n such that the n th term of the series is less than 1.

D6**Use sequences and series in modelling.**

Only assessed at A-level

Teaching guidance

Students should be able to answer questions set within a context, eg compound interest.

Examples

- 1 A ball is dropped from a height of 1 metre above the ground.

Each time it hits the ground it bounces to a height of $\frac{3}{4}$ the distance it fell before the bounce.

- (a) Show that the distance travelled by the ball between the first and second bounce is 1.5 metres.
 - (b) Find the total distance travelled by the ball after it is dropped.
- 2 A maths graduate begins a job with a starting salary of £26 000. She has been promised an annual pay rise of 5% of the previous year's salary.
- (a) How much should she expect to earn in her third year?
 - (b) If she stays in the same job for seven years, how much would she earn in total over this time?

Note: A-level students should be encouraged to apply their mathematical thinking to problem solving and modelling from the first day of teaching.

E

Trigonometry

E1

Understand and use the definitions of sine, cosine and tangent for all arguments; the sine and cosine rules; the area of a triangle in the form $\frac{1}{2}ab \sin C$


Work with radian measure, including use for arc length and area of sector.

Assessed at AS and A-level

Teaching guidance

Students should:


- know and be able to apply the following rules:
in any triangle ABC
- area of triangle $\frac{1}{2}ab \sin C$
- sine rule $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
- cosine rule $a^2 = b^2 + c^2 - 2bc \cos A$



- be aware of the ambiguous case that can arise from the use of the sine rule.

Examples


- 1 The triangle ABC , shown in the diagram, is such that $BC = 6$ cm, $AC = 5$ cm and $AB = 4$ cm. The angle BAC is θ .



- (a) Use the cosine rule to show that $\cos \theta = \frac{1}{8}$
- (b) Find the exact value of $\sin \theta$
- (c) Find the area of the triangle ABC .
- 2 Angle θ is such that $\sin \theta = \frac{1}{7}$ and $0 < \theta < 90^\circ$

Find the exact values of $\cos \theta$ and $\tan \theta$

- 3 The diagram shows a triangle ABC .



The lengths of the sides AC and BC are 5 cm and 6 cm respectively.

The area of triangle ABC is 12.5 cm^2 , and angle ACB is obtuse.

- (a) Find the size of angle ACB , giving your answer to the nearest 0.1°
- (b) Find the length of AB , giving your answer to two significant figures.

Only assessed at A-level


Teaching guidance

Students should:

- understand and be able to use radian measure.
- know that $2\pi \text{ rad} = 360^\circ$
- know and be able to use $l = r\theta$, $A = \frac{1}{2}r^2\theta$

Examples

- 1 The diagram shows a sector of OAB of a circle with centre O and radius $r \text{ cm}$.



The angle AOB is 1.25 radians. The perimeter of the sector is 39 cm.

- Show that $r = 12$
- Calculate the area of the sector OAB .

E2

Understand and use the standard small angle approximations of sine, cosine and tangent

$$\sin \theta \approx \theta$$

$$\cos \theta \approx 1 - \frac{\theta^2}{2}$$

$$\tan \theta \approx \theta$$

where θ is in radians.

Only assessed at A-level

Teaching guidance

Students should be able to:

- understand and use the standard small angle approximations of sine, cosine and tangent when differentiating sine or cosine from first principles.
- use standard small angle approximations to deduce approximations for other functions.

Examples

- 1 (a) Show, using a small angle approximation, that $\sec x \approx \frac{2}{2-x^2}$
 - (b) Hence, find the first two terms of the binomial expansion for $\sec x$
 - (c) Using your binomial expansion find an approximate value for $\sec 0.1$, giving your answer to 5 decimal places.
- 2 (a) Using a compound angle identity write down an expression for $\sin(x+h)$
 - (b) Using small angle approximations for $\sin(h)$ and $\cos(h)$, and your answer to part (a), find

$$\lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin(x)}{h}$$

E3

Understand and use the sine, cosine and tangent functions; their graphs, symmetries and periodicity.

Know and use exact values of \sin and \cos for

$0, \frac{\pi}{6}, \frac{\pi}{4}, \frac{\pi}{3}, \frac{\pi}{2}, \pi$ and multiples thereof, and exact values

of \tan for $0, \frac{\pi}{6}, \frac{\pi}{4}, \frac{\pi}{3}, \pi$ and multiples thereof.

Assessed at AS and A-level

Teaching guidance


Students should be able to:

- understand and use vertical asymptotes of a tangent graph.
- carry out simple transformations of the graphs of the sine, cosine and tangent functions.

Note: radians will not be required at AS.

Examples

- 1 (a) On the axes given below, sketch the graph of $y = \tan x$, for $0^\circ \leq x \leq 360^\circ$



- (b) Solve the equation $\tan x = -1$, giving all the values of x in the interval $0^\circ \leq x \leq 360^\circ$

- 2 Find the two solutions of the equation $\sin x = \sin \sqrt{41}$ for $0^\circ \leq x \leq 360^\circ$

Give your answers in exact form.

Teaching guidance

Only assessed at A-level

Students should:

- understand and be able to use radians.
- understand the phrase 'exact value' and use their calculators appropriately (many calculators will give the required exact form).
- understand the difference between showing a result and simply evaluating on a calculator (to avoid questions that require trivial calculator results, students may be asked to show particular results).

Examples

- 1 Given that $\sin a = \frac{1}{3}$ and $0 < a < \frac{\pi}{2}$, find the exact value of $\sin\left(a + \frac{\pi}{6}\right)$
- 2 (a) Sketch the graph of $y = \cos x$ in the interval $0 \leq x \leq 2\pi$
(b) State the values of the intercepts with the coordinate axes.

E4

Understand and use the definitions of secant, cosecant and cotangent and of arcsin, arccos and arctan; their relationships to sine, cosine and tangent; understanding their graphs; their ranges and domains.

Only assessed at A-level

Teaching guidance

Students should:

- know and be able to use the following functions and their graphs:

Secant	Cosecant	Cotangent
$\sec x = \frac{1}{\cos x}$	$\operatorname{cosec} x = \frac{1}{\sin x}$	$\cot x = \frac{1}{\tan x} = \frac{\cos x}{\sin x}$

- be aware of both notations, ie $\arcsin x$ or $\sin^{-1}x$;
- know the domains and ranges.
- know that:

$$-\frac{\pi}{2} \leq \sin^{-1}x \leq \frac{\pi}{2}$$


$$0 \leq \cos^{-1}x \leq \pi$$

$$-\frac{\pi}{2} < \tan^{-1}x < \frac{\pi}{2}$$

- understand how to sketch the graphs by reflecting relevant sections of the trigonometric graphs in the line $y = x$
- be able to apply simple transformations to graphs of all these functions.

Examples


- 1 The diagram shows the curve $y = \cos^{-1}x$ for $-1 \leq x \leq 1$



Write down the exact coordinates of points A and B .

- 2 Sketch the curve with equation $y = \operatorname{cosec} x$ for $0 < x < \pi$


- 3 (a) The sketch shows the graph of $y = \sin^{-1}x$



Write down the coordinates of the points P and Q , the end points of the graph.



- (b) Sketch the graph of $y = \sin^{-1}(x-1)$
- 4 Solve the equation $\sec x = 5$, giving all the values of x in the interval $0 \leq x \leq 2\pi$ in radians to two decimal places.

- 5 (a) The diagram shows the graph of $y = \sec x$ for $0^\circ \leq x \leq 360^\circ$



- (i) The point A on the curve is where $x = 0$. State the y -coordinate of A.
 - (ii) Sketch the graph of $y = \sec(2x) + 2$ for $0^\circ \leq x \leq 360^\circ$
- (b) Solve the equation $\sec x = 2$ giving all the values of x in degrees in the interval $0^\circ \leq x \leq 360^\circ$

- 6 (a) Sketch the graph of $y = \cos^{-1} x$, where y is in radians on the first axes below. State the coordinates of the end points of the graph.
- (b) Sketch the graph of $y = \pi - \cos^{-1} x$, where y is in radians on the second axes below. State the coordinates of the end points of the graph.



E5

Understand and use $\tan \theta = \frac{\sin \theta}{\cos \theta}$

Understand and use $\sin^2 \theta + \cos^2 \theta = 1$;
 $\sec^2 \theta = 1 + \tan^2 \theta$ and $\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- use $\tan \theta = \frac{\sin \theta}{\cos \theta}$ to solve equations or find exact values.
- use $\sin^2 \theta + \cos^2 \theta = 1$ to solve equations or find exact values.

Examples

- (a) Given that $6 \tan \theta \sin \theta = 5$, show that $6 \cos^2 \theta + 5 \cos \theta - 6 = 0$

(b) Hence solve the equation $6 \tan (3x) \sin (3x) = 5$, giving all values of x to the nearest degree in the interval $0^\circ \leq x \leq 180^\circ$
- It is given that $(\tan \theta + 1)(\tan^2 \theta - 3) = 0$

(a) Find the possible values of $\tan \theta$

(b) Hence solve the equation $(\tan \theta + 1)(\sin^2 \theta - 3 \cos^2 \theta) = 0$, giving all solutions for θ , in degrees, in the interval $0^\circ \leq \theta \leq 180^\circ$
- Given that $\sin \theta = \frac{2}{7}$ and θ is obtuse, find:

(a) the exact value of $\cos \theta$

(b) the exact value of $\tan \theta$

Only assessed at A-level

Teaching guidance

Students should be able to:

- use $\sec^2\theta = 1 + \tan^2\theta$ and $\operatorname{cosec}^2\theta = 1 + \cot^2\theta$ to solve equations or find exact values.
- use identities to perform integration, eg $\int \tan^2 x \, dx$
- prove identities.

Examples

- 1 (a) By using a suitable trigonometrical identity, solve the equation

$$\tan^2\theta = 3(3 - \sec\theta)$$

giving all solutions to the nearest 0.1° in the interval $0^\circ < \theta < 360^\circ$

- (b) Hence solve the equation

$$\tan^2(4x - 10^\circ) = 3[3 - \sec(4x - 10^\circ)]$$

giving all solutions to the nearest 0.1° in the interval $0^\circ < x < 90^\circ$

E6

Understand and use double angle formulae; use of formulae for $\sin(A \pm B)$, $\cos(A \pm B)$ and $\tan(A \pm B)$; understand geometrical proofs of these formulae.

Understand and use expressions for $a\cos\theta + b\sin\theta$ in the equivalent forms of $r\cos(\theta \pm \alpha)$ or $r\sin(\theta \pm \alpha)$

Only assessed at A-level

Teaching guidance

Students should be able to:

- understand and use the addition formulae:

$$\sin(A \pm B) = \sin(A)\cos(B) \pm \cos(A)\sin(B)$$

$$\cos(A \pm B) = \cos(A)\cos(B) \mp \sin(A)\sin(B)$$

$$\tan(A \pm B) = \frac{\tan(A) \pm \tan(B)}{1 \mp \tan(A)\tan(B)}$$

- understand how these formulae can be used to derive the double angle formulae:

$$\sin(2A) = 2\sin(A)\cos(A)$$

$$\cos(2A) = \cos^2(A) - \sin^2(A)$$

$$= 2\cos^2(A) - 1$$

$$= 1 - 2\sin^2(A)$$

- use double angle formulae to solve equations and integration.
- use $r\cos(\theta \pm \alpha)$ or $r\sin(\theta \pm \alpha)$ to solve equations or describe features of the resulting wave function, eg maximum or minimum, amplitude, etc.

Notes: The identities may be written using either \equiv or $=$ and either will be accepted in exams.

Examples

- 1 (a) Express $\cos 2x$ in the form $a\cos^2 x + b$, where a and b are constants.

- (b) Hence show that $\int_0^{\frac{\pi}{2}} \cos^2 x \, dx = \frac{\pi}{4}$, where a is an integer.

- 2 It is given that $3\cos\theta - 2\sin\theta = R\cos(\theta + \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$
- Find the value of R .
 - Show that $\alpha \approx 33.7^\circ$
 - Hence write down the maximum value of $3\cos\theta - 2\sin\theta$ and find a positive value of θ at which this maximum value occurs.
- 3 The polynomial $f(x)$ is defined by $f(x) = 4x^3 - 11x - 3$
- Use the Factor Theorem to show that $(2x + 3)$ is a factor of $f(x)$
 - Write $f(x)$ in the form $(2x + 3)(ax^2 + bx + c)$
 - Show that the equation $2\cos 2\theta \sin\theta + 9\sin\theta + 3 = 0$ can be written as $4x^3 - 11x - 3 = 0$, where $x = \sin\theta$
 - Hence find all solutions of the equation $2\cos 2\theta \sin\theta + 9\sin\theta + 3 = 0$ in the interval $0^\circ < \theta < 360^\circ$, giving your solutions to the nearest degree.
- 4
- Show that $\cot x - \sin 2x = \cot x \cos 2x$ for $0^\circ < x < 180^\circ$
 - Hence, or otherwise, solve the equation $\cot x - \sin 2x = 0$
in the interval $0^\circ < x < 180^\circ$
- 5 Angle α is acute and $\cos\alpha = \frac{3}{5}$. Angle β is obtuse and $\sin\beta = \frac{1}{2}$.
- Find the value of $\tan\alpha$ as a fraction.
 - Find the value of $\tan\beta$ in surd form.
 - Hence show that

$$\tan(\alpha + \beta) = \frac{m\sqrt{3} - n}{n\sqrt{3} + m}$$
 where m and n are integers.

E7

Solve simple trigonometric equations in a given interval, including quadratic equations in **sin**, **cos** and **tan** and equations involving multiples of the unknown angle.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand and solve simple trigonometric equations.
- answer questions that require them to give solutions in degrees.

Examples

- 1 Solve the equation $\sin(\theta - 30^\circ) = 0.7$, giving your answers to the nearest 0.1° in the interval $0^\circ \leq \theta \leq 360^\circ$
- 2 Write down the two solutions of the equation $\tan x = \tan 61^\circ$ in the interval $0^\circ \leq x \leq 360^\circ$
- 3 Solve the equation $\sin 2x = \sin 48^\circ$, giving the values of x in the interval $0^\circ \leq x \leq 360^\circ$
- 4 (a) Given that

$$\frac{\cos^2 x + 4\sin^2 x}{1 - \sin^2 x} = 7$$

show that

$$\tan^2 x = \frac{3}{2}$$

- (b) Hence solve the equation

$$\frac{\cos^2 2\theta + 4\sin^2 2\theta}{1 - \sin^2 2\theta} = 7$$

in the interval

$$0^\circ \leq \theta \leq 180^\circ$$

giving your values of θ to the nearest degree.

Only assessed at A-level

Teaching guidance

Students should be able to:

- understand and solve trigonometric equations using formulae from section E6,
eg $\sec\left(2x + \frac{\pi}{3}\right) = \sqrt{2}$
- answer questions that require them to give solutions in radians.

Examples

1 Solve the equation $\tan \frac{1}{2}x = 3$ in the interval $0 < x < 4\pi$, giving your answers in radians to three significant figures.

2 Solve the equation $\cos \theta (\sin \theta - 3 \cos \theta) = 0$ in the interval $0 < \theta < 2\pi$, giving your answers in radians to three significant figures.

3 It is given that $2 \operatorname{cosec}^2 x = 5 - 5 \cot x$.

(a) Show that the equation $2 \operatorname{cosec}^2 x = 5 - 5 \cot x$ can be written in the form

$$2 \cot^2 x + 5 \cot x - 3 = 0$$

(b) Hence show that $\tan x = 2$ or $\tan x = -\frac{1}{3}$

(c) Hence, or otherwise, solve the equation $2 \operatorname{cosec}^2 x = 5 - 5 \cot x$, giving all values of x in radians to one decimal place in the interval $-\pi \leq x \leq \pi$

3 (a) Solve the equation $\sec x = 5$, giving all the values of x in the interval $0 \leq x \leq 2\pi$ in radians to two decimal places.

(b) Show that the equation $\tan^2 x = 3 \sec x + 9$ can be written in the form

$$\sec^2 x - 3 \sec x - 10 = 0$$

(c) Solve the equation $\tan^2 x = 3 \sec x + 9$, giving all the values of x in the interval $0 \leq x \leq 2\pi$ in radians to two decimal places.

E8

Construct proofs involving trigonometric functions and identities.

Only assessed at A-level

Teaching guidance

Students should understand and be able to use the following:

$$\sin^2\theta + \cos^2\theta \equiv 1$$

$$\sec^2\theta \equiv 1 + \tan^2\theta$$

$$\operatorname{cosec}^2\theta \equiv 1 + \cot^2\theta$$

Note: unless stated in the question, the above may be quoted without proof.

Examples

1 Prove the identity $(\tan x + \cot x)^2 \equiv \sec^2 x + \operatorname{cosec}^2 x$

2 Prove the identity $\frac{1 - \sin x}{\cos x} + \frac{\cos x}{1 - \sin x} \equiv 2 \sec x$

3 Prove the identity $\frac{\sec^2 x}{(\sec x + 1)(\sec x - 1)} \equiv \operatorname{cosec}^2 x$

E9

Use trigonometric functions to solve problems in context, including problems involving vectors, kinematics and forces.

Only assessed at A-level

Teaching guidance

Students should be able to:


- solve problems using any of the techniques from sections E1 to E8 on their own or in combination.
- select for themselves the appropriate technique for solving a problem.

Examples

- 1 A crane is lowering a heavy crate down a mine shaft when the crate scrapes the side of the mine shaft, twists and becomes stuck.


The mine shaft has a width of 2 metres and the crate is 1.8 metres tall by 1.2 metres wide.

The angle between the wall of the mine shaft and the side of the crate is θ , as shown in the diagram.




- (a) Show that $9\sin \theta + 6\cos \theta = 10$
- (b) Hence find the value of θ

- 2 Two vectors \overrightarrow{AB} and \overrightarrow{BC} have magnitude 7.6 units and 5.8 units respectively and the obtuse angle between the vectors is 118° , as shown in the diagram:



Find the magnitude of the vector \overrightarrow{AC} .

- 3 A particle is in equilibrium under the action of four horizontal forces of magnitudes 5 N, 8 N, P N and Q N, as shown in the diagram:



- (a) Show that $P = 9$.
- (b) Find the value of Q .

Note: A-level students should be encouraged to apply their mathematical thinking to problem solving and modelling from the start of the course.

F

Exponentials and logarithms

F1

Know and use the function a^x and its graph, where a is positive.

Know and use the function e^x and its graph.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- sketch and use simple transformations of the function a^x .
- sketch and use simple transformations of the function e^x .

Note: at A-level, students should also be able to sketch and use **a combination** of simple transformations of the functions a^x and e^x .

Examples

- Sketch the graph of $y = 3^x$, stating the coordinates of the point where the graph crosses the y -axis.
 - Describe a single geometrical transformation that maps the graph of $y = 3^x$:
 - onto the graph of $y = 3^{2x}$
 - onto the graph of $y = 3^{x+1}$

- The curve $y = 3 \times 12^x$ is translated by the vector $\begin{pmatrix} 1 \\ p \end{pmatrix}$ to give the curve $y = f(x)$.

Given that the curve $y = f(x)$ passes through the origin $(0, 0)$, find the value of the constant p .

- Sketch the graph of $y = 9^x$, indicating the value of the intercept on the y -axis.
 - The curve $y = 9^x$ is reflected in the y -axis to give the curve with equation $y = f(x)$. Write down an expression for $f(x)$.

F2

Know that the gradient of e^{kx} is equal to ke^{kx} and hence understand why the exponential model is suitable in many applications.

Assessed at AS and A-level

Teaching guidance

At AS, students should know that the gradient is proportional to the value of the function

At AS, students are **not** expected to differentiate functions involving e^{kx} .

Examples

- 1 Find the gradient of the curve $y = e^{2x}$ at the point where $x = 2$.

Circle your answer.

$2e$

$2e^2$

e^4

$2e^4$

- 2 A model for the growth of a colony of bacteria is $P = 500e^{\frac{1}{8}t}$, where P is the number of bacteria after t minutes.

What is the initial rate of growth of the bacteria?

Only assessed at A-level

Teaching guidance

Students should understand and be able to use exponential functions at a more in-depth level.

Example

- 1 A biologist is investigating the growth of a population of a species of rodent. The biologist proposes the model

$$N = \frac{500}{1 + 9e^{-\frac{t}{8}}}$$

for the number of rodents, N , in the population t weeks after the start of the investigation.

- (a) Show that the rate of growth, $\frac{dN}{dt}$, is given by

$$\frac{dN}{dt} = \frac{N}{4000}(500 - N)$$

- (b) The maximum growth occurs after T weeks. Find the value of T .

F3

Know and use the definition of $\log_a x$ as the inverse of a^x , where a is positive and $x \geq 0$.

Know and use the function $\ln x$ and its graph.

Know and use $\ln x$ as the inverse function of e^x .

Assessed at AS and A-level

Teaching guidance

Students should:

- understand and be able to use the equivalence of $y = a^x \Leftrightarrow \log_a y = x$ and $y = e^x \Leftrightarrow \ln y = x$
- know that the graph of $y = \ln x$ is a reflection in the line $y = x$ of the graph of $y = e^x$
- be able to perform simple single transformations of the functions $y = e^x$ and $y = \ln x$
- be able to manipulate logs and exponentials if required within the solution to a problem.

Examples

- Sketch the graph of $y = 2\ln x$ indicating any points where the curve crosses the coordinate axes.
 - The graph of $y = 2\ln x$ can be transformed into the graph of $y = 1 + 2\ln x$ by means of a translation. Write down the vector of the translation.

- If $A = B^n$, which of the following is true?

Circle your answer.

$$n = \log_B A$$

$$n = \log_A B$$

$$B = \log_n A$$

$$B = \log_A n$$

- Given that $\log_a b = c$, express b in terms of a and c .

Only assessed at A-level

Teaching guidance

Students should be able to:


- sketch a combination of transformations of the functions $y = e^x$ and $y = \ln x$.
- use a combination of transformations of the functions $y = e^x$ and $y = \ln x$.
- use the terms domain and range in problems using the functions $y = e^x$ and $y = \ln x$.

1 A function is defined by $f(x) = 2e^{3x} - 1$ for all real values of x .

(a) Find the range of f .

(b) Show that $f^{-1}(x) = \frac{1}{3} \ln\left(\frac{x+1}{2}\right)$

2 The curve with equation $y = f(x)$, where $f(x) = \ln(2x-3)$, $x > \frac{3}{2}$, is sketched below.



The inverse of f is f^{-1}

- (a) Find $f^{-1}(x)$
- (b) State the range of f^{-1}
- (c) Sketch the curve with equation $y = f^{-1}(x)$, indicating the coordinates of the point where the curve intersects the y -axis.

F4

Understand and use the laws of logarithms:

$$\log_a x + \log_a y = \log_a (xy);$$

$$\log_a x - \log_a y = \log_a \left(\frac{x}{y} \right);$$

$$k \log_a x = \log_a x^k$$

(including, for example, $k = -1$ and $k = -\frac{1}{2}$).

Assessed at AS and A-level

Teaching guidance

Students should:

- know, understand and be able to use the above laws.
- know that $\log_a a = 1$ and $\log_a 1 = 0$ for $a > 0$

Examples

- (a) Solve the equation $3\log_a x = \log_a 8$

(b) Show that $3\log_a 6 - \log_a 8 = \log_a 27$
- (a) Given that $\log_a x = 2\log_a 6 - \log_a 3$, show that $x = 12$

(b) Given that $\log_a y = \log_a 5 + 7$, express y in terms of a , giving your answer in a form not involving logarithms.

Only assessed at A-level

Example

- The curve $y = 3^x$ intersects the line $y = x + 3$ at the point where $x = \alpha$

(a) Show that α lies between 0.5 and 1.5.

(b) Show that the equation $3^x = x + 3$ can be arranged into the form

$$x = \frac{\ln(x+3)}{\ln 3}$$

F5**Solve equations of the form $a^x = b$.**

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- solve equations of the form $a^x = b$, including $e^x = b$

Examples

- 1 The line $y = 85$ intersects the curve $y = 6^{3x}$ at the point A . By using logarithms, find the x -coordinate of A , giving your answer to three decimal places.
- 2 The line $y = 21$ intersects the curve $y = 3(2^x + 1)$ at the point P .
 - (a) Show that the x -coordinate of P satisfies the equation $2^x = 6$
 - (b) Use logarithms to find the x -coordinate of P , giving your answer to three significant figures.
- 3 Given that $e^{-2x} = 3$, find the exact value of x .

F6

Use logarithmic graphs to estimate parameters in relationships of the form $y = ax^n$ and $y = kb^x$, given data for x and y .

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- reduce a non-linear relationship to linear form.
- plot graphs from given data, drawing a line of best fit by eye and using it to calculate gradients and intercepts as estimates for unknown constants.


Note: this is an essential skill in A-level Sciences and there is an ideal opportunity here to link to real data. You can search for Power laws for relationships of the form $y = ax^n$ and exponential laws for those of the form $y = kb^x$

Examples

- 1 The variables y and x are related by an equation of the form $y = ax^n$ where a and n are constants.

Let $Y = \log_{10} y$ and $X = \log_{10} x$.

- (a) Show that there is a linear relationship between Y and X .
- (b) The graph of Y against X is shown in the diagram.



Find the value of n and the value of a .

- 2 The variables x and y are known to be related by an equation in the form $y = ab^x$ where a and b are constants.


The following approximate values of x and y have been found.

x	1	2	3	5
y	3.84	6.14	9.82	15.8

- (a) Complete the table, showing values of x and Y , where $Y = \log_{10} y$. Give each value of Y to three decimal places.

x	1	2	3	4
Y	0.584			

- (b) Show that, if $y = ab^x$, then x and Y must satisfy an equation of the form $Y = mx + c$.
- (c) Draw a graph relating x and Y .



- (d) Hence find estimates for the values of a and b .

F7

Understand and use exponential growth and decay; use in modelling (examples may include the use of e in continuous compound interest, radioactive decay, drug concentration decay, exponential growth as a model for population growth); consideration of limitations and refinements of exponential models.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- use given conditions to determine the values of unknown constant terms in equations of the forms $y = Ae^{bx} + C$ or $P = Ak^t + C$
- form and use exponential equations to make predictions.

Examples

- 1 On 1 January 1900, a sculpture was valued at £80.

When the sculpture was sold on 1 January 1956, its value was £5000.

The value, £ V , of the sculpture is modelled by the formula $V = Ak^t$, where t is the time in years since 1 January 1900 and A and k are constants.

- Write down the value of A .
- Show that $k = 1.07664$ to five decimal places.
- Use this model to:
 - show that the value of the sculpture on 1 January 2006 will be greater than £200 000.
 - find the year in which the value of the sculpture will first exceed £800 000.

- 2 A biologist is researching the growth of a certain species of hamster. She proposes that the length, x cm, of a hamster t days after its birth is given by $x = 15 - 12e^{-\frac{t}{14}}$
- (a) Use this model to find:
- (i) the length of a hamster when it is born.
 - (ii) the length of hamster after 14 days, giving your answer to three significant figures.
- (b) (i) Show that the time for a hamster to grow to 10 cm in length is given by
- $$t = 14 \ln \left(\frac{a}{b} \right)$$
- where a and b are integers.
- (ii) Find this time to the nearest day.
- 3 The concentration, $C \text{ mg l}^{-1}$, of a particular drug in a patient's bloodstream t hours after it has been administered is given by the formula $C = C_0 e^{-0.2t}$
- (a) Initially a patient is given a dose of 5 mg l^{-1}
- (i) Write down the value of C_0
 - (ii) Find the concentration of the drug 3 hours after it is administered.
- (b) The drug becomes ineffective when the concentration drops below 2 mg l^{-1}
- How long does it take for the drug to become ineffective? Give your answer in hours to three significant figures.

Note: within every set of exam papers at AS, 20% of the assessment must address AO3. At A-level, 25% of assessment addresses AO3.

Examples of modelling should be introduced to students at an early stage of teaching so that they can build confidence in the use of models and in the interpretation of the outputs from mathematical models. Models will not always be given to students. Students will sometimes be required to translate a situation in context into a mathematical model. Teachers should be mindful of both OT2 and OT3 because many assessment items will be set in the context of these overarching themes, which address problem-solving and modelling.

G

Differentiation

G1

Understand and use the derivative of $f(x)$ as the gradient of the tangent to the graph of $y = f(x)$ at a general point (x, y) ; the gradient of the tangent as a limit; interpretation as a rate of change; sketching the gradient function for a given curve; second derivatives; differentiation from first principles for small positive integer powers of x and for $\sin x$ and $\cos x$.

Understand and use the second derivative as the rate of change of gradient; connection to convex and concave sections of curves and points of inflection.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- recognise and use the notations $f'(x)$, $\frac{dy}{dx}$, $\frac{d}{dx}(f(x))$
- use $f''(x)$ or $\frac{d^2y}{dx^2}$ to determine the nature of a stationary point.

Examples

- 1 A model helicopter takes off from a point O at time $t = 0$ and moves vertically so its height, y cm, above O after time t seconds is given by:

$$y = \frac{1}{4}t^4 - 26t^2 + 96t, \quad 0 \leq t \leq 4$$

(a) Find:

(i) $\frac{dy}{dt}$

(ii) $\frac{d^2y}{dt^2}$

- (b) Verify that y has a stationary value when $t = 2$ and determine whether this stationary value is a maximum value or a minimum value.
- (c) Find the rate of change of y with respect to t when $t = 1$
- (d) Determine whether the height of the helicopter above O is increasing or decreasing at the instant when $t = 3$
- (e) Determine whether the speed of the helicopter is increasing or decreasing at the instant when $t = 3$

- 2 The volume, $V \text{ m}^3$, of water in a tank at time t seconds is given by

$$V = \frac{1}{3}t^6 - 2t^4 + 3t^2 \text{ for } t \geq 0$$


(a) Find:

(i) $\frac{dV}{dt}$


(ii) $\frac{d^2V}{dt^2}$

- (b) Find the rate of change of the volume of water in the tank, in $\text{m}^3 \text{ s}^{-1}$, when $t = 2$

- 3 The graph of $y = f(x)$ is shown below.



Sketch the graph of $y = f'(x)$ on the axes below.



- 4 A curve has equation $y = x^3 - 12x$

The point A on the curve has coordinates $(2, -16)$

The point B on the curve has x -coordinate $2 + h$

- Show that the gradient of the line AB is $6h + h^2$
- Explain how the result of part (a) can be used to show that A is a stationary point on the curve.

Only assessed at A-level

Teaching guidance

Students should:

- know that at points of inflection $f''(x) = 0$, but that the converse is not necessarily true.
- know that for concave and convex sections over the interval $[a, b]$ the following holds:
 - a twice-differentiable function is concave $\Leftrightarrow f''(x) \leq 0$ for all $x \in [a, b]$.
 - a twice-differentiable function is convex $\Leftrightarrow f''(x) \geq 0$ for all $x \in [a, b]$.

Examples

- 1 A curve is given by the equation $f(x) = x^3 - 3x^2 + 1$
Find the range of values of x for which the curve is concave.

- 2 The point $A\left(\frac{\pi}{3}, \frac{1}{2}\right)$ is on the curve $y = \cos x$

The point B on the curve has x -coordinate $\frac{\pi}{3} + h$

- (a) Show that the gradient of the line AB can be written as

$$\frac{\cos h - \sqrt{3} \sin h - 1}{2h}$$

- (b) Using the result from part (a) and small angle approximations explain how the gradient of the curve, $y = \cos x$, at the point A can be found.

G2

Differentiate x^n , for rational values of n , and related constant multiples, sums and differences.

Differentiate e^{kx} and a^{kx} , $\sin kx$, $\cos kx$ and $\tan kx$ related sums, differences and constant multiples. Understand and use the derivative of $\ln x$.

Assessed at AS and A-level

Teaching guidance

Students should know and be able to use the following:

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\frac{d}{dx}(Ax^n) = Anx^{n-1}$$

$$\frac{d}{dx}(Ax^n + Bx^m) = Anx^{n-1} + Bmx^{m-1}$$

Example

- 1 A model car moves so that its distance, x centimetres, from a fixed point O after time t seconds is given by

$$x = \frac{1}{2}t^4 - 20t^2 + 66t$$

Find:

(a) $\frac{dx}{dt}$

(b) $\frac{d^2x}{dt^2}$

Only assessed at A-level

Teaching guidance

Students should know and be able to use the following:

$f(x)$	$f'(x)$
e^{kx}	ke^{kx}
$\ln x$	$\frac{1}{x}$
$\sin kx$	$k \cos kx$
$\cos kx$	$-k \sin kx$

The following are given in the Formulae booklet:

a^{kx}	$(k \ln a)a^{kx}$
$\tan kx$	$k \sec^2 x$

Examples

1 Find $\frac{dy}{dx}$ when $y = e^{3x} + \ln x$

2 A curve has the equation $y = e^{2x} - 10e^2 + 12x$

(a) Find $\frac{dy}{dx}$

(b) Find $\frac{d^2y}{dx^2}$

3 Find $\frac{dy}{dx}$ when $y = \tan 3x$

G3

Apply differentiation to find gradients, tangents and normals, maxima and minima and stationary points, points of inflection.

Identify where functions are increasing or decreasing.

Assessed at AS and A-level

Teaching guidance

Students should:

- understand and be able to use the fact that at a stationary point, $\frac{dy}{dx} = 0$
- describe a stationary point as a (local) maximum or minimum
- know that:

At a maximum $\frac{dy}{dx} = 0$ and $\frac{d^2y}{dx^2} < 0$

At a minimum $\frac{dy}{dx} = 0$ and $\frac{d^2y}{dx^2} > 0$

Note:

the case $\frac{d^2y}{dx^2} = 0$ will not be tested at AS

- use $m_1 \times m_2 = -1$ for gradients of tangents and normal.
- be able to answer questions set in the form of a practical problem where a function of a single variable has to be optimised
- be able to show that a function is increasing or decreasing, by showing $\frac{dy}{dx} > 0$ or $\frac{dy}{dx} < 0$ respectively.

Examples

1 The curve with equation $y = x^4 - 32x + 5$ has a single stationary point, M .


(a) Find $\frac{dy}{dx}$

(b) Hence find the coordinates of M .

(c) (i) Find $\frac{d^2y}{dx^2}$

(ii) Hence, or otherwise, determine whether M is a maximum or minimum point.

2 The diagram shows a block of wood in the shape of a prism with a triangular cross-section. The end faces are right-angled triangles with sides of lengths $3x$ cm, $4x$ cm and $5x$ cm, as shown in the diagram.



The total surface area of the five faces is 144 cm^2 .

(a) (i) Show that $xy + x^2 = 12$.

(ii) Hence show that the volume of the block, $V \text{ cm}^3$, is given by $V = 72x - 6x^3$


(b) (i) Find $\frac{dV}{dx}$

(ii) Show that V has a stationary value when $x = 2$

(c) (i) Find $\frac{d^2V}{dx^2}$

(ii) Hence determine whether V has a maximum value or a minimum value when $x = 2$

- 3 A curve, drawn from the origin O , crosses the x -axis at the point $P(4, 0)$.
The normal to the curve at P meets the y -axis at the point Q , as shown in the diagram.



The curve, defined for $x \geq 0$, has the equation $y = 4x^{\frac{1}{2}} - x^{\frac{3}{2}}$

- (a) Find $\frac{dy}{dx}$
 - (b) Show that the gradient of the curve at $P(4, 0)$ is -2
 - (c) Find an equation of the normal to the curve at $P(4, 0)$
 - (d) Find the y -coordinate of Q and hence find the area of triangle OPQ .
- 4 A curve has the equation $y = e^{-2x} + 6x$
- (a) Find the exact values of the coordinates of the stationary point of the curve.
 - (b) Determine the nature of the stationary point.

Only assessed at A-level

Teaching guidance

Students should:

- know that a point of inflection is where a curve changes from convex to concave or vice versa.
- understand that $\frac{d^2y}{dx^2} = 0$ at a point of inflection, but this is not, on its own, sufficient to show the existence of a point of inflection, eg $y = x^4$ has a minimum at $x = 0$ (not a point of inflection) and $\frac{d^2y}{dx^2} = 0$ when $x = 0$

Example

- 1 A curve has the equation $y = e^x + e^{-x}$

Show that any points of inflection satisfy $e^{2x} = -1$

G4

Differentiate using the product rule, the quotient rule and the chain rule, including problems involving connected rates of change and inverse functions.

Only assessed at A-Level

Teaching guidance

Students should:

- know and be able to use the following rules:

Product rule	$\frac{d}{dx}[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$
Quotient rule (this is given in the Formulae booklet)	$\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$
Chain rule	<p>If $y = f(u)$ and $u = g(x)$, so that $y = f(g(x))$ then</p> $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$
Inverses	$\frac{dx}{dy} = \frac{1}{\frac{dy}{dx}}$

- be able to use product, quotient or chain rules for differentiating $\tan x$, $\sec x$, $\operatorname{cosec} x$ and $\cot x$

Examples

1 (a) Find $\frac{dy}{dx}$ when:

(i) $y = (4x^2 + 3x + 2)^{10}$

(ii) $y = x^2 \tan x$

(b) (i) Find $\frac{dy}{dx}$ when $x = 2y^3 + \ln y$

(ii) Hence find an equation of the tangent to the curve $x = 2y^3 + \ln y$ at the point (2, 1).

2 (a) Find $\frac{dy}{dx}$ when:

(i) $y = (2x^2 - 5x + 1)^{20}$

(ii) $y = x \cos x$

(b) Given that

$$y = \frac{x^3}{x-2}$$

show that

$$\frac{dy}{dx} = \frac{kx^2(x-3)}{(x-2)^2}$$

where k is a positive integer.

3 A bucket is being used to catch water dripping from a leaking classroom ceiling, at a constant rate of 0.5 cm^3 per minute.

The volume, $V \text{ cm}^3$, of water in the bucket is given by

$$V = \frac{\pi h}{3}(h^2 + 3h + 300)$$

Where h is the depth of water in the bucket.

Find the rate of change of h in cm per minute when the depth of water in the bucket is 10 cm.

4 The function f is defined by $f(x) = \sin x$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$

Find $\frac{d}{dx}(f^{-1}(x))$ in terms of x .

G5

Differentiate simple functions and relations defined implicitly or parametrically, for first derivative only.

Only assessed at A-Level

Teaching guidance

Students should be able to:

- use the chain rule with parametric equations: eg for $y = e^t + e^{-t}$ and $x = e^t - e^{-t}$

$$\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx} \quad \text{or} \quad \frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$$
- use the chain rule with parametric equations to find stationary points, equations of tangents and normals, but **not** to find points of inflection **nor** to test for concavity.

Examples

- 1 A curve is defined by the parametric equations

$$x = 1 + 2t, \quad y = 1 - 4t^2$$

(a) (i) Find $\frac{dx}{dt}$ and $\frac{dy}{dt}$

(ii) Hence find $\frac{dy}{dx}$ in terms of t .

(b) Find an equation of the normal to the curve at the point where $t = 1$.

(c) Find a Cartesian equation of the curve.

- 2 A curve is defined by the equation

$$y^2 - xy + 3x^2 - 5 = 0$$

(a) Find the y -coordinates of the two points on the curve where $x = 1$.

(b) (i) Show that $\frac{dy}{dx} = \frac{y-6x}{2y-x}$

(ii) Find the gradient of the curve at each of the points where $x = 1$.

(iii) Show that, at the two stationary points on the curve, $33x^2 - 5 = 0$

- 3 A curve is defined by the parametric equations $x = 2\cos\theta$, $y = 3\sin 2\theta$

(a) (i) Show that

$$\frac{dy}{dx} = a\sin\theta + b\operatorname{cosec}\theta$$

where a and b are integers.

(ii) Find the gradient of the normal to the curve at the point where $\theta = \frac{\pi}{6}$

(b) Show that the Cartesian equation of the curve can be expressed as

$$y^2 = px^2(4 - x^2)$$

where p is a rational number.

- 4 A curve is defined by the equation $9x^2 - 6xy + 4y^2 = 3$

Find the coordinates of the two stationary points of this curve.

- 5 (a) A curve is defined by the equation $x^2 - y^2 = 8$

(i) Show that at any point (p, q) on the curve, where

$$q \neq 0$$

the gradient of the curve is given by

$$\frac{dy}{dx} = \frac{p}{q}$$

(ii) Show that the tangents at the points (p, q) and $(p, -q)$ intersect on the x -axis.

(b) Show that

$$x = t + \frac{2}{t}, \quad y = t - \frac{2}{t}$$

are parametric equations of the curve

$$x^2 - y^2 = 8$$

G6

Construct simple differential equations in pure mathematics and in context. (Contexts may include kinematics, population growth and modelling the relationship between price and demand.)

Only assessed at A-level

Teaching guidance

Students should understand language associated with proportionality and rates of change.

Examples

- 1 A water tank has a height of 2 metres. The depth of the water in the tank is h metres at time t minutes after water begins to enter the tank. The rate at which the depth of water in the tank increases is proportional to the difference between the height of the tank and the depth of the water.

Write down a differential equation in the variables h and t and a positive constant k .

- 2 A biologist is investigating the growth of a population of a species of rodent. The biologist proposes the model

$$N = \frac{500}{1 + 9e^{-\frac{t}{8}}}$$

for the number of rodents, N , in the population, t weeks after the start date of the investigation.

Use this model to answer the following questions.

- (a) Show that the rate of growth

$$\frac{dN}{dt}$$

is given by

$$\frac{dN}{dt} = \frac{N}{4000}(500 - N)$$

- (b) The maximum rate of growth occurs after T weeks. Find the value of T .

- 3 A giant snowball is melting. The snowball can be modelled as a sphere whose surface area is decreasing at a constant rate with respect to time. The surface area of the sphere is $A \text{ cm}^2$ at a time t days after it begins to melt.

Write down a differential equation in terms of the variables A and t and a constant k , where $k > 0$, to model the surface area of the melting snowball.

- 4 The number of fish in a lake is decreasing. After t years, there are x fish in the lake. The rate of decrease of the number of fish is proportional to the number of fish currently in the lake.
- (a) Formulate a differential equation, in the variables x and t and a constant of proportionality k , where $k > 0$, to model the rate at which the number of fish in the lake is decreasing.
- (b) At a certain time, there were 20 000 fish in the lake and the rate of decrease was 500 fish per year. Find the value of k .

Note: students should be prepared to encounter unstructured questions that address AO3. One of the aims of this new A-level is to encourage students to use their mathematical skills and techniques to solve challenging problems which require the students to decide on the solution strategy. From the early stages of teaching, the specification encourages students to take decisions and devise their own strategies when faced with an unstructured question.

H

Integration

H1

Know and use the Fundamental Theorem of Calculus.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand that differentiation is the ‘reverse’ of integration and vice versa
- use

$$\int_b^a f(x) \, dx = F(b) - F(a)$$

where

$$\frac{d}{dx}(F(x)) = f(x)$$

Note: the maximum level of difficulty for questions at AS requires students to use an integrand, $f(x)$, where $f(x)$ is the sum of terms of the form ax^n where n is rational and $n \neq -1$

Examples

1 Find $\int_{-2}^0 (x^3 - x + 6) \, dx$

Note: a question asked in this style will not be worth many marks as calculators can evaluate the answer directly and the use of technology is to be encouraged.

2 Find $\int_0^2 \sqrt{x} \, dx$ giving your answer in the form $\frac{a\sqrt{2}}{b}$

Note: a question of this style will be worth more marks, as permitted calculators will not give the exact form required.

Only assessed at A-level

Example

1 Find the exact value of $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \sin x \, dx$.

H2

Integrate x^n (excluding $n = -1$), and related sums, differences and constant multiples.

Integrate e^{kx} , $\frac{1}{x}$, $\sin kx$, $\cos kx$ and related sums, differences and constant multiples.

Assessed at AS and A-level

Teaching guidance

Students should:

- know that:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$$

and

$$\int ax^n + bx^m dx = a \int x^n dx + b \int x^m dx$$

- understand integration as the reverse of differentiation and should include a constant of integration when finding an indefinite integral.

Examples

- 1 (a) The expression $\left(1 - \frac{1}{x^2}\right)^3$ can be written in the form $1 + \frac{p}{x^2} + \frac{q}{x^4} - \frac{1}{x^6}$.

Find the values of the integers p and q .

- (b) Hence find $\int \left(1 - \frac{1}{x^2}\right)^3 dx$.

- 2 Find $\int \left(1 + 3x^{\frac{1}{2}} + x^{\frac{3}{2}}\right) dx$.

Only assessed at A-level

Teaching guidance

Students should know and be able to use the following:

$f(x)$	$\int f(x) dx$
e^{kx}	$\frac{1}{k}e^{kx} + c$
$\frac{1}{x}$	$\ln x + c$
$\sin kx$	$-\frac{1}{k}\cos kx + c$
$\cos kx$	$\frac{1}{k}\sin kx + c$

Examples

- 1 Find $\int 2e^{4x} + \frac{1}{2x} dx$
- 2 (a) Using the product rule, find $\frac{dy}{dx}$ when $y = x \ln x$
 (b) Hence find $\int \ln x dx$
- 3 Find $\int 4\sin 2x - 12\cos 3x dx$

H3

Evaluate definite integrals; use a definite integral to find the area under a curve and the area between two curves.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand and use the fact that for a function, f , where $f(x) \geq 0$ for $a \leq x \leq b$ the area between the x -axis the curve $y = f(x)$ and the lines $x = a$ and $x = b$ is given by


$$\text{area} = \int_a^b f(x) dx$$

- understand that for areas lying **below** the x -axis the definite integral will give the negative of the required value.
- Find areas between curves and straight lines.

Note: at AS curves will be entirely above or entirely below the x -axis for the required areas.

Examples

- The curve with equation $y = x^4 - 8x + 9$ is sketched below.




The point $(2, 9)$ lies on the curve.


(a) Find $\int_0^2 (x^4 - 8x + 9) dx$

- (b) Hence find the area of the shaded region bounded by the curve and the line $y = 9$

- 2 The curve C with equation $y = x^3 + x - 10$, sketched below, crosses the x -axis at the point $Q(2, 0)$



- Find the gradient of the curve C at the point Q .
 - Hence find an equation of the tangent to the curve C at the point Q .
 - Find $\int (x^3 + x - 10) dx$.
 - Hence find the area of the shaded region bounded by the curve C and the coordinate axes.
- 3 The curve with equation $y = x^3 + 4x - 5$ is sketched below.



The curve cuts the x -axis at the point $A(1, 0)$ and the point $B(2, 11)$ lies on the curve.

Find the area of the shaded region bounded by the curve and the line AB .


Only assessed at A-level

Teaching guidance

Students should understand that extra care must be taken when finding areas where curves cross the x -axis, or where there are two curves, and the total area of more than one region is required.

Example

- 1 The two curves, $y = \sin x$ and $y = \sin 2x$, for $0 \leq x \leq \pi$, are shown on the graph.



- (a) Find the exact coordinates of the point of intersection A.
- (b) Find the exact value of the area of the shaded region.

H4

Understand and use integration as the limit of a sum.

Only assessed at A-level

Teaching guidance

Students should be able to:


- understand that the area under a curve can be approximated using rectangles. The limit as the number of rectangles is increased is equal to a definite integral.
- recognise and use notation such as $\lim_{n \rightarrow \infty} \sum_{i=1}^n y_i \delta x = \int_a^b y \, dx$

Notes: the idea here is to link to work at GCSE using the trapezium rule and have a semi-formal understanding of the ideas of Riemann sums, although such vocabulary is not required. Students should comment on whether an approximation would give an over- or underestimate or if it cannot be decided, linking this to the ideas of increasing and decreasing functions.

The ideas here can be investigated using interactive applets such as intmath.com/integration/riemann-sums.php This is an example of how the use of technology should permeate the course.

Example

- 1 The area under the graph of $y = x^3 - 3x + 3$ between the lines $x = 1$ and $x = 2$ is to be estimated by calculating the areas of the 8 rectangles, of equal width, δx , as shown in the diagram.



- (a) What is the value of δx ?
- (b) The area of the smallest bar can be found using the calculation $y_1 \times \delta x$, where y_1 is the height of the bar.

Calculate the total area of all the bars

$$\sum_{i=1}^8 y_i \delta x$$

- (c) A more accurate estimate of the area under the curve can be found by using more bars of smaller width.

Calculate

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n y_i \delta x$$

H5

Carry out simple cases of integration by substitution and integration by parts; understand these methods as the inverse processes of the chain and product rules respectively.

(Integration by substitution includes finding a suitable substitution and is limited to cases where one substitution will lead to a function which can be integrated; integration by parts includes more than one application of the method but excludes reduction formulae.)

Only assessed at A-level

Teaching guidance

Students should be able to:

- understand the simplest cases of substitution such as:

$$\int e^{ax+b} dx$$

$$\int \sin(ax+b) dx$$

$$\int \frac{1}{ax+b} dx$$

Note: it is likely students will learn these as standard integrals rather than use substitution each time.

- spot integrals of the form

$$\int f'(x) \cdot [f(x)]^n dx$$

and integrate directly

- use $\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$

to perform integration by parts, limited to a maximum of two applications of the method.

Examples

1 Find $\int x^2 \sin 2x dx$

- 2 (a) Given that

$$y = \frac{\sin \theta}{\cos \theta}$$

use the quotient rule to show that

$$\frac{dy}{d\theta} = \sec^2 \theta$$

- (b) Given that

$$x = \sin \theta$$

show that

$$\frac{x}{\sqrt{1-x^2}} = \tan \theta$$

- (c) Use the substitution $x = \sin \theta$ to find

$$\int \frac{1}{(1-x^2)^{\frac{3}{2}}} dx$$

giving your answer in terms of x

- 3 (a) (i) Find $\int \ln x \, dx$

- (ii) Find $\int (\ln x)^2 \, dx$

- (b) Use the substitution $u = \sqrt{x}$ to find the exact value of $\int_1^4 \frac{1}{x + \sqrt{x}} \, dx$


- 4 Using integration by substitution, or otherwise, find $\int x^2 (x^3 + 2)^7 \, dx$

- 5 Find $\int \frac{x^2 + 2x}{2x^3 + 6x^2 + 1} \, dx$

Note: it is more likely that questions will be asked like this, with no scaffolding.

- 6 Find $\int \frac{\sin x}{\cos x} \, dx$

- 7 The curve with parametric equations $y = 1 - \cos \theta$ and $x = \cos \theta$, $0 \leq \theta \leq 2\pi$ is shown below.




Find the area enclosed between the curve and the x -axis.

- 8 (a) Given that $y = x^{-2} \ln x$, show that $\frac{dy}{dx} = \frac{1 - 2 \ln x}{x^3}$

(b) Using integration by parts, find $\int x^{-2} \ln x \, dx$

(c) The sketch shows the graph of $y = x - 2 \ln x$



Using the answer to part (a), find, in terms of e , the x -coordinate of the stationary point A .

H6

Integrate using partial fractions that are linear in the denominator.

Only assessed at A-level

Teaching guidance

Students should be able to answer questions that require the simplification of a more complicated fraction, which leads to an integrable expression or to partial fractions that can be integrated.

Examples

1 (a) (i) Express

$$\frac{5x-6}{x(x-3)}$$

in the form

$$\frac{A}{x} + \frac{B}{x-3}$$

(ii) Find $\int \frac{5x-6}{x(x-3)} dx$.

(b) (i) Given that

$$4x^3 + 5x - 2 = (2x+1)(2x^2 + px + q) + r$$

find the values of the consonants p , q and r .

(ii) Find $\int \frac{4x^3 + 5x - 2}{2x+1} dx$

2 (a) (i) Express

$$\frac{5-8x}{(2+x)(1-3x)}$$

in the form

$$\frac{A}{2+x} + \frac{B}{1-3x}$$

where A and B are integers.

(ii) Hence show that

$$\int_{-1}^0 \frac{5-8x}{(2+x)(1-3x)} dx = p \ln 2$$

where p is rational.

(b) (i) Given that

$$\frac{9-18x-6x^2}{2-5x-3x^2}$$

can be written as

$$C + \frac{5-8x}{2-5x-3x^2}$$

find the value of C .

(ii) Hence find the exact value of the area of the region bounded by the curve

$$y = \frac{9-18x-6x^2}{2-5x-3x^2}, \text{ the } x\text{-axis and the lines } x = -1 \text{ and } x = 0.$$

You may assume that $y > 0$ when $-1 \leq x \leq 0$

3 (a) Given that

$$\frac{9x^2-6x+5}{(3x-1)(x-1)}$$

can be written in the form

$$3 + \frac{A}{3x-1} + \frac{B}{x-1}$$

where A and B are integers, find the values of A and B .

(b) Hence, or otherwise, find $\int \frac{9x^2-6x+5}{(3x-1)(x-1)} dx$

H7

Evaluate the analytical solution of simple first order differential equations with separable variables, including finding particular solutions.

(Separation of variables may require factorisation involving a common factor.)

Only assessed at A-level

Teaching guidance

Students should be able to carry out any of the techniques of integration included in sections H1 to H6.

Examples

- 1 Solve the differential equation

$$\frac{dy}{dx} = 6xy^2$$

given that $y = 1$ when $x = 2$.

Give your answer in the form $y = f(x)$.

- 2 Solve the differential equation

$$\frac{dy}{dx} = \frac{1}{y} \cos\left(\frac{x}{3}\right)$$

given that

$$y = 1 \text{ when } x = \frac{\pi}{2}$$

Write your answer in the form $y^2 = f(x)$.

- 3 (a) (i) Solve the differential equation

$$\frac{dy}{dt} = y \sin t$$

to obtain y in terms of t .

- (ii) Given that $y = 50$ when $t = \pi$, show that $y = 50e^{-(1+\cos t)}$

- 4 Find the general solution of the differential equation

$$\frac{dy}{dx} = xy + 2x$$

Give your answer in the form $y = f(x)$.

- 5 Solve the differential equation

$$\frac{dy}{dx} = \frac{x\sqrt{x^2 + 3}}{e^{2y}}$$

given that $y = 0$ when $x = 1$.

Give your answer in the form $y = f(x)$

H8

Interpret the solution of a differential equation in the context of solving a problem, including identifying limitations of the solution; includes links to kinematics.

Only assessed at A-level

Teaching guidance

Students should be able to solve differential equations they have set up themselves.

Examples

- 1 A car's value **depreciates** at a rate which is proportional to its value, £ V , at time t months from when it was new.
 - (a) Write down a differential equation in terms of the variables V and t and a constant k , where $k > 0$, to model the value of the car.
 - (b) Solve your differential equation to show that $V = Ad^t$ where $d = e^{-k}$
 - (c) The value of the car when new was £12 499 and 36 months later its value was £7000.
Find the values of A and d .

- 2 The platform of a theme park ride oscillates vertically. For the first 75 seconds of the ride

$$\frac{dx}{dt} = \frac{t \cos\left(\frac{\pi}{4}t\right)}{32x}$$

where x metres is the height of the platform above the ground after time t seconds.

At $t = 0$, the height of the platform above the ground is 4 metres.

Find the height of the platform after 45 seconds, giving your answer to the nearest centimetre.

Note: the construction of, solution of, and interpretation of the solutions relating to differential equations in context should be encouraged.

I

Numerical methods

I1

Locate roots of $f(x) = 0$ by considering changes of sign of $f(x)$ in an interval of x on which $f(x)$ is sufficiently well-behaved.

Understand how change of sign methods can fail.

Only assessed at A-level

Teaching guidance

Students should:

- be able to answer questions where roots of equations in the form $g(x) = h(x)$ can be located by rearranging to give $f(x) = g(x) - h(x) = 0$
- know that discontinuities can cause failure but also know that the converse is not true.

Examples

- 1 The curve $y = 3^x$ intersects the line $y = x + 3$ at the point where $x = \alpha$.

Show that α lies between 0.5 and 1.5.

- 2 Show that the equation

$$x^3 - 6x + 1 = 0$$

has a root α , where

$$2 < \alpha < 3$$

- 3 When searching for a root to the equation $f(x) = 0$, a student correctly evaluates $f(1) = 4$ and $f(2) = 3$

Given that f is a continuous function, which one of the following could not be true?

Circle the correct answer.

There are
no roots
for $1 < x < 2$

There is
one root
for $1 < x < 2$

There are
two roots
for $1 < x < 2$

There are
four roots
for $1 < x < 2$

12

Solve equations approximately using simple iterative methods; draw associated cobweb and staircase diagrams.

Solve equations using the Newton-Raphson method and other recurrence relations of the form $x_{n+1} = g(x_n)$

Understand how such methods can fail.

Only assessed at A-level

Teaching guidance

Students should be able to:

- answer questions that require them to rearrange an equation into an iterative form.
- draw staircase and cobweb diagrams, to illustrate the iteration, on printed graphs.
- demonstrate divergence on a diagram.
- understand the conditions when the Newton-Raphson method may fail.

Notes: this is another opportunity to use technology. This GeoGebra interactive allows you to investigate cobweb diagrams: [geogebra.org/m/XvjM7Xnv](https://www.geogebra.org/m/XvjM7Xnv)

Students should expect to use calculators to perform iterative operations.

Examples

- 1 (a) The equation $e^{-x} - 2\sqrt{x} = 0$ has a single root, α .
Show that α lies between 3 and 4.

- (b) Use the recurrence relation


$$x_{n+1} = (2 - e^{-x_n})^2$$

With

$$x_1 = 3.5$$

to find x_2 and x_3 giving your answers to three decimal places.

- (c) The diagram shows parts of the graphs of $y = (2 - e^{-x})^2$ and $y = x$, and a position of x_1 .



On the diagram, draw a staircase or cobweb diagram to show how convergence takes place, indicating the positions of x_2 and x_3 on the x -axis.

- 2 The equation $24x^3 + 36x^2 + 18x - 5 = 0$ has one real root, α .

- (a) Show that α lies in the interval $0.1 < x < 0.2$
- (b) Taking $x_1 = 0.2$ as a first approximation to α , use the Newton-Raphson method to find a second approximation, x_2 , to α . Give your answer to four decimal places.

- 3 The equation $x^3 - x^2 - 2 = 0$ has a single root between 1 and 2.

Starting with $x_1 = 1$ which of the following iterative formulae will not converge on the root?

Circle your answer.

$$x_{n+1} = \sqrt{\frac{(x_n)^2 + 2}{x_n}}$$

$$x_{n+1} = \sqrt[3]{(x_n)^2 + 2}$$

$$x_{n+1} = \frac{2}{(x_n)^2} + 1$$

$$x_{n+1} = \frac{(x_n)^3 - 2}{x_n}$$

- 4 A curve has equation $y = x^3 - 3x + 3$.

(a) Show that the curve intersects the x -axis at the point $(\alpha, 0)$ where $-3 < \alpha < -2$.

(b) A student attempts to find α using the Newton-Raphson method with $x_1 = -1$.

Explain why the student's method fails.

- 5 Use the iteration

$$x_{n+1} = \frac{\ln(x_n + 3)}{\ln 3}$$

with

$$x_1 = 0.5$$

to find x_3 to two significant figures.

I3

Understand and use numerical integration of functions, including the use of the trapezium rule and estimating the approximate area under a curve and limits that it must lie between.

Only assessed at A-level

Teaching guidance

Students should be able to:

- understand and use the term 'ordinate'.
- use graphical determination to find whether an approximation over- or under-estimates the area.
- improve an approximation by increasing the number of ordinates or strips used.

This topic links to H4 and can again be usefully explored using interactives.

Examples


- 1 (a) Use the trapezium rule with five ordinates (four strips) to find an approximate value for

$$\int_0^4 \frac{1}{x^2 + 1} dx$$

giving your answer to four significant figures.

- (b) State how you could obtain a better approximation to the value of the integral using the trapezium rule.

- 2 The diagram shows a sketch of the curve $y = 2^{4x}$



The curve intersects the y -axis at the point A .

- (a) Find the value of the y -coordinate of A .
- (b) Use the trapezium rule with six ordinates (five strips) to find an approximate value for

$$\int_0^1 2^{4x} dx$$

giving your answer to two decimal places.

- 3 The trapezium rule is used, with six ordinates, to find an estimate for the value of

$$\int_0^5 f(x) dx$$

Given that f is an increasing function and $f''(x) > 0$, explain what would happen to the value obtained by the trapezium rule if the number of ordinates were increased.

14

Use numerical methods to solve problems in context.

Only assessed at A-level

Teaching guidance

Students should be able to:

- use a numerical method to solve an equation or to calculate an area resulting from a problem that has its origin within pure mathematics or from a context such as exponential growth or kinematics.
- demonstrate that they can apply the correct numerical method in full. Numerical solutions obtained directly from calculator functions are unlikely to achieve full credit.

Examples

- 1 A particle travels in a straight line with its velocity, $v \text{ m s}^{-1}$, at time, t seconds, given by

$$v = 10e^{\frac{t}{5}} \sin t$$

Use the trapezium rule, with 6 ordinates, to estimate the distance moved by the particle in its first 5 seconds of motion.

- 2 A particle is projected vertically upwards so that its height, h metres, above the ground after time t seconds is given by

$$h = 20t - 9.8t^2 + 2.5e^{-t}$$

- (a) How high above the ground was the point where the particle was initially launched?
- (b) Given that the particle is in the air for between 1.5 and 2.5 seconds, use the Newton-Raphson method to find the time when the particle hits the ground. Give your answer to two significant figures.

Note: teachers should remind students of OT2.4: many mathematical problems cannot be solved analytically, but numerical methods permit a solution to a required level of accuracy. This is a subtle and very important point: students are often comfortable with what they consider trial and error methods; we teach them analytical methods that are generally more efficient, but now we are saying that sometimes there are problems that require numerical methods.

J

Vectors

J1

Use vectors in two dimensions and in three dimensions.

Assessed at AS and A-level

Teaching guidance

Students should:

- become familiar with both column vectors and \mathbf{i} , \mathbf{j} notation, where \mathbf{i} and \mathbf{j} are unit vectors in perpendicular directions. Unless specified otherwise, students may use either notation in their solutions.

Note: questions may be set without context as pure vector questions, or in context, for example, using vectors to represent velocities, forces etc.

- know that vectors may be used to describe translations of graphs.

Notes: our standard notation will be to use round brackets for column vectors, but square brackets are also acceptable. When writing vectors using single lower case letters, students may underline or overline, but no particular notation will be required.

Examples

- 1 The points A and B have coordinates $(-2, 3)$ and $(4, -1)$ respectively.

Write down the vector \overrightarrow{AB} in the form $a\mathbf{i} + b\mathbf{j}$.

- 2 A force, \mathbf{F} , acts on a particle of mass 2 kg.

Given that $\mathbf{F} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ N, find the acceleration of the particle.

Only assessed at A-level

Teaching guidance

Students should become familiar with both column vectors and \mathbf{i} , \mathbf{j} , \mathbf{k} notation, where \mathbf{i} , \mathbf{j} and \mathbf{k} are unit vectors in mutually perpendicular directions in a right-handed coordinate system. Unless specified otherwise, students may use either notation in their solutions.

Example

- 1 The points A and B have coordinates $(-2, 3, 1)$ and $(4, -1, -2)$ respectively.

Write down the vector \overrightarrow{AB} in the form $a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$.

J2

Calculate the magnitude and direction of a vector and convert between component form and magnitude/direction form.

Assessed at AS and A-level

Teaching guidance

Students should be able to answer questions using vectors in two dimensions.

Examples

- 1 Calculate the angle between the vector \mathbf{j} and the vector $2\mathbf{i} + 3\mathbf{j}$.
- 2 A cricket ball is hit at ground level on a horizontal surface.
It initially moves at 26 m s^{-1} at an angle of 50° above the horizontal.
 - (a) Find the horizontal component of the ball's initial velocity, giving your answer to two significant figures.
 - (b) Write down the initial velocity of the ball as a vector in the form $a\mathbf{i} + b\mathbf{j}$
- 3 Two forces, $\mathbf{P} = (6\mathbf{i} - 3\mathbf{j}) \text{ N}$ and $\mathbf{Q} = (3\mathbf{i} + 15\mathbf{j}) \text{ N}$, act on a particle.
The unit vectors \mathbf{i} and \mathbf{j} are perpendicular.
 - (a) State the resultant of \mathbf{P} and \mathbf{Q} .
 - (b) Calculate the magnitude of the resultant of \mathbf{P} and \mathbf{Q} .
 - (c) Calculate the direction of the resultant of \mathbf{P} and \mathbf{Q} .

Note: when answering part (c), students would be expected to be specific about where they are measuring their direction from. This can be done with a clear diagram or described as 53.1° from the \mathbf{i} direction, for example.

4 A ship is sailing with a constant velocity of $\begin{pmatrix} 5 \\ -7 \end{pmatrix}$ km/h.

- (a) Find the speed of the ship.
- (b) Find the direction in which the ship is sailing, giving your answer as a 3-figure bearing to the nearest degree.

Only assessed at A-level

Teaching guidance

Students should be able to answer questions using vectors in two or three dimensions.

Example

1 The points A and B have coordinates $(2, 5, 1)$ and $(4, 1, -2)$ respectively.

- (a) Find the vector \overrightarrow{AB} .
- (b) Find the length AB .

J3

Add vectors diagrammatically and perform the algebraic operations of vector addition and multiplication by scalars, and understand their geometrical interpretations.

Assessed at AS and A-level

Teaching guidance


Students should be able to:

- prove that two vectors are parallel to each other by showing that one is a multiple of the other.
- recall the conditions for collinearity.
- understand that a vector diagram can be used to find resultants. This could, for example, be in the context of force.

Note: unless specifically stated in a question, students can choose **any** appropriate method to solve vector problems.

Examples

1




$$\overrightarrow{OP} = \overrightarrow{PQ} = \mathbf{a}$$

$$\overrightarrow{OM} = \overrightarrow{MR} = \mathbf{b}$$

The point T is on the line PR such that $PT:TR = 1:2$

- Find \overrightarrow{PT} in terms of \mathbf{a} and \mathbf{b}
- Prove that M , T and Q are collinear.

- 2 The diagram shows three forces which act in the same plane and are in equilibrium.



- (a) Find F
- (b) Find α

Only assessed at A-level

Example

- 1 The points A , B and C have coordinates $(3, 1, -6)$, $(5, -2, 0)$ and $(8, -4, -6)$ respectively.

- (a) Show that the vector \overrightarrow{AC} is given by $\overrightarrow{AC} = n \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}$, where n is an integer.
- (b) The point D is such that $ABCD$ is a rhombus.
 - (i) Find the coordinates of D .
 - (ii) Calculate the side length of the rhombus.

J4

Understand and use position vectors; calculate the distance between two points represented by position vectors.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand and use the result $\overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA}$.
- use position vectors in both a pure context and in mechanics questions, for example in kinematics.

Example

- 1 The points A and B have position vectors $\overrightarrow{OA} = 5\mathbf{i} - 2\mathbf{j}$ and $\overrightarrow{OB} = -\mathbf{i} + 3\mathbf{j}$

Find the distance between the points A and B .

Only assessed at A-level

Examples

- 1 A particle is initially at the point A , which has position vector $13.6\mathbf{i}$ metres, with respect to an origin O .

At the point A , the particle has velocity $(6.0\mathbf{i} + 2.4\mathbf{j}) \text{ m s}^{-1}$ and in its subsequent motion, it has a constant acceleration of $(-0.80\mathbf{i} + 0.10\mathbf{j}) \text{ m s}^{-2}$.

The unit vectors \mathbf{i} and \mathbf{j} are directed east and north respectively.

Find an expression for the position vector of the particle, with respect to the origin O , t seconds after it leaves A .

- 2 The positions of two particles, A and B , at a time t , are given by $\mathbf{r}_A = 3 \cos(\pi t)\mathbf{i} + (2t + 1)\mathbf{j}$ and $\mathbf{r}_B = 2 \sin(\pi t)\mathbf{i} + (2 - t)\mathbf{j}$ respectively.

Find the distance between the two particles when $t = 4$.

Note: in mechanics questions where measured quantities have been given to varying degrees of accuracy students should give their final answer to an appropriate degree of accuracy. For example, if some measured lengths are given to two significant figures, but other measured lengths are given to three significant figures the final answer should be given to two significant figures.

J5

Use vectors to solve problems in pure mathematics and in context, including forces and kinematics.

Assessed at AS and A-level

Teaching guidance

Students should be able to use vectors in 2 dimensions to solve problems.

Note: at AS, questions on kinematics will **not** involve vectors but questions related to forces may involve 2-D vectors.

Example

1 Three forces $\mathbf{F}_1 = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$ N, $\mathbf{F}_2 = \begin{pmatrix} 7 \\ 3 \end{pmatrix}$ N and $\mathbf{F}_3 = \begin{pmatrix} a \\ b \end{pmatrix}$ N act at a point and are in equilibrium.

Find a and b .

Only assessed at A-level

Teaching guidance

Students should be able to:

- use vectors in up to **3** dimensions to solve problems.
- use vectors to solve problems related to kinematics in up to **2** dimensions, including projectile motion.

Example

1 Three points A , B and C are collinear with $\overrightarrow{OA} = \begin{pmatrix} 1 \\ 5 \\ -2 \end{pmatrix}$ and $\overrightarrow{OB} = \begin{pmatrix} 3 \\ -1 \\ 1 \end{pmatrix}$.

Given that $AC = 2AB$ find the coordinates of the two possible positions of point C .

For sections K to O students must use a calculator that can find:

- exact and cumulative probabilities in the binomial distribution
- probabilities in a normal distribution.

Teachers are advised to teach as much as possible within the context of the large data set that is current within the specification.

K

Statistical sampling

K1

Understand the terms ‘population’ and ‘sample’.

Use samples to make informal inferences about the population.

Understand and use sampling techniques, including simple random sampling and opportunity sampling.

Select or critique sampling techniques in the context of solving a statistical problem, including understanding that different samples can lead to different conclusions about the population.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- identify a population and understand, and explain, the meaning of the terms ‘parameter’ and ‘statistic’.
- understand that information from a sample can be used to make inferences about the population it was taken from.
- recognise and explain how to use sampling techniques, to include simple random sampling, systematic sampling, opportunity sampling, stratified sampling, quota sampling and cluster sampling.
- understand that the use of prior information can make the sample more representative of the population (eg stratification).
- decide which sampling method to use to overcome practical problems when sampling.
- discuss the advantages and disadvantages of each sampling method within a given context.

Examples

- 1 A large number of spectators attend a football match at a ground in England. One of the stands contains 1390 seats, numbered from 1 to 1390.
- (a) (i) Describe how random numbers could be used to select a random sample of 80 of these seats.
- (ii) The ground authority wishes to carry out a survey into spectators' opinions of the catering facilities at the ground. It proposes to ask the spectators occupying the 80 seats selected in part (a) (i) to complete a questionnaire.
- Suggest **two** practical difficulties that might be encountered in carrying out this proposal.
- (b) A market research company is employed to investigate the amount of interest taken in sport by the population of the United Kingdom. James, a new employee of the company, suggests that interviewers should be stationed outside each exit from the ground with instructions to interview every 100th person leaving the ground after the match.
- (i) State the name given to this type of sampling.
- (ii) Give **one** reason why the results of the investigation suggested by James might be biased.
- (iii) Suggest **two** practical difficulties, other than those which you mentioned in part (a) (ii), which might be encountered in carrying out these interviews.

- 2 A fan club has 2076 members who are divided geographically into eight branches. The club's committee wishes to seek members' views on where to hold the next annual meeting. A sample of 100 members is to be obtained and their views sought.

The following suggestions are made as to how to choose the sample.

Suggestion A Members are selected from all eight branches. The number of members from each branch is proportional to the size of the branch. The branch secretaries are asked to choose the appropriate number of members from their branches in any convenient way.

Suggestion B Two of the branches are selected at random. Fifty members from each of these branches are selected at random.

Suggestion C Members are selected by a random process from each branch. The number of members from each branch is proportional to the size of the branch.

Suggestion D Members are selected by the chair of the committee, from the branch she is a member of, at the next local meeting.

- (a) For **each** of the four suggestions:
- (i) name the method of sampling.
 - (ii) either state that each member is equally likely to be included in the sample, or explain why this is not the case.
- (b) (i) State, giving a reason, which of the four methods is preferable from a statistical point of view.
- (ii) Give a reason why Suggestion A might be preferred to Suggestion C.

- 3 Emile wants to investigate the proportion of the population who watch soap operas on television.

One day, he asks each of the students in his drama class: "Did you watch a soap opera on television last night?"

- (a) Name the type of sampling that Emile is using.
- (b) Comment on the use of the proportion of students who watched a soap opera last night, obtained from Emile's sample, to be representative of the proportion of the population who watch soap operas on television.

Note: part (b) of this question does not specify exactly what a student is required to include in their response. This leaves it to the student to decide on their own response. Possible responses to this may be along the following lines:

Emile's opportunity sample is reflecting only a very restricted sample of the population, namely only students/drama students in his class at one school/college in the country.

Not every member of the population under investigation is equally likely to be included in the sample so the results are likely to contain bias.

It is unlikely that the proportion of students who watched a soap opera last night will be representative of the proportion of the population who watch soap operas.

- 4 Packets of a particular type of sweet are known to have a mean of 100 grams. The number of sweets in a packet is approximately 30 and the sweets come in any one of five flavours. The weights of 50 packets are taken and the mean is found to be 98.3 grams.

From the above passage, identify:

- (a) a population.
 - (b) a parameter.
 - (c) a sample.
 - (d) a qualitative variable.
 - (e) a continuous variable.
 - (f) a discrete variable.
- 5 Hermione wants to investigate the proportion of people in her school who enjoy watching horror movies.
- She decides to go from table to table in the school refectory and to ask the question 'do you enjoy watching horror movies?' to each table as a group.
- She will record the number of 'yes' responses and the total number of students for each table.
- Hermione assumes that each student will respond independently to her question.
- (a) Comment on the assumption that each student will respond independently to her question.
 - (b) Suggest how Hermione could change her method of collecting data to improve the reliability of her data.

Note: this example asks students to consider how more reliable data may be achieved. This may be achieved, for example, by getting each student to record their response on a piece of paper and folding this before putting it into a box so that no other student sees their response (anonymity being more likely to elicit a genuine response rather than one influenced by other students on the table). Students may also suggest that respondents could be asked to complete their responses online where there is no mention of who is conducting the investigation and that this may then reduce possible bias in responses given etc.

L

Data presentation and interpretation

L1

Interpret diagrams for single-variable data, including understanding that area in a histogram represents frequency.

Connect to probability distributions.

Assessed at AS and A-level

Teaching guidance


Students should be able to:

- interpret box and whisker plots, cumulative frequency curves and histograms.
Note: students will **not** be expected to construct these diagrams.
- use diagrams to find probabilities of given events.

Note: when studying this topic, students can use data from the large data set and process it using software such as GeoGebra ([geogebra.org](https://www.geogebra.org)) which has a wealth of materials available to download.

Examples

- 1 Rehana wishes to catch a train from her local station to the city centre. There are two local taxi companies; Blue Star and Green Star. In the past Rehana has telephoned both companies. The times, in minutes, that she has to wait between telephoning and the arrival of a taxi are summarised in the box and whisker plot.




- (a) Compare briefly the waiting times for the two taxi companies.
- (b) Giving a reason for your choice, advise Rehana on which company to telephone if, in order to catch the next train, she needs the taxi to arrive within:
- (i) 5 minutes.
 - (ii) 25 minutes.

- 2 Sally's Safaris is a holiday company which organises adventurous holidays. The ages, in years, of the customers who booked holidays in the year 2002 are summarised in the table below.

Age	Frequency
15-24	35
25-29	29
30-34	20
35-44	27
45-59	30
60-79	28

Sally drew the histogram below to illustrate the data. Unfortunately, both coordinates of the point marked R have been plotted incorrectly.

State the correct coordinates of the point R .



L2

Interpret scatter diagrams and regression lines for bivariate data, including recognition of scatter diagrams which include distinct sections of the population (calculations involving regression lines are excluded).
Understand informal interpretation of correlation.
Understand that correlation does not imply causation.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- interpret a scatter diagram, to include visual recognition of outliers (as stated in section L4).
- recognise and name positive, negative or no correlation as types of correlation.
- recognise and name strong, moderate or weak correlation as strengths of correlation.
- understand that just because a correlation exists it does not necessarily mean that a causality is present. This is spurious correlation.
- state and use the fact that $-1 \leq r \leq 1$
- interpret, in context, correlation by considering a scatter diagram or a given value for r .

Note: students will **not** be required to calculate or evaluate a correlation coefficient. Students will **not** be required to calculate the equation of a regression line or to make predictions using calculations using the equation of a regression line.

Students will not be expected to plot scatter diagrams, but should be encourage to use software to plot data from the LDS on scatter diagrams.

Examples

- 1 Dr Hanna has a special clinic for her older patients. She asked a medical student, Lenny, to select a random sample of 25 of her male patients, aged between 55 and 65 years, and from their clinical records, to list their heights, weights and waist measurements.


Lenny was then asked to calculate three values of the product moment correlation coefficient based upon his collected data. His results were:

- (a) 0.365 between height and waist measurement
- (b) 1.16 between height and weight
- (c) -0.583 between weight and waist measurement.


For each of Lenny's three calculated values, state whether the value is definitely correct, probably correct, probably incorrect or definitely incorrect.

2 For each of the diagrams below, comment on the type of correlation illustrated.


(a)



(b)



(c)



Note: students are left to decide that (b) illustrates both moderate and negative correlation and that (c) can be interpreted to show overall negative correlations but that there are apparently two separate groups of data that both illustrate strong and positive correlation.

L3

Interpret measures of central tendency and variation, extending to standard deviation.

Calculate standard deviation, including from summary statistics.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- find the mean, median, mode, range, quartiles and interquartile range from data given in graphical or tabular form.
- interpret values of the mean, median and mode.
- calculate standard deviation (or variance) using a calculator or from summary statistics of the form $\sum x$, $\sum x^2$ or $\sum (x - \bar{x})^2$

Notes: it is advisable for students to know whether to divide by n or $(n - 1)$ when calculating either the variance of a population or an estimate for the population from a sample data. However, either divisor will be accepted **unless** a question specifically requests an unbiased estimate of a population variance.

For small data sets, the positions of the median and quartiles are usually given by

$\frac{n+1}{4}$, $\frac{n+1}{2}$, $\frac{3(n+1)}{4}$ and it will often be convenient to ensure 1 is a multiple of 4.

However, the quartiles are more relevant to large sets of data and here it is usually more convenient to replace $n+1$ by n .

Examples

- 1 The runs scored by a cricketer in 11 innings during the 2006 season were as follows.

47 63 0 28 40 51 a 77 0 13 35

The exact value of a was unknown but it was greater than 100.

- Calculate the median and the interquartile range of these 11 values.
- Give a reason why, for these 11 values:
 - the mode is **not** an appropriate measure of average.
 - the range is **not** an appropriate measure of spread.

- 2 The weight of fat in a digestive biscuit is known to be normally distributed.

Pat conducted an experiment in which she measured the weight of fat, x grams, in each of a random sample of 10 digestive biscuits, with the following results:

$$\sum x = 31.9 \text{ and } \sum (x - \bar{x})^2 = 1.849$$

Use this information to calculate estimates of the mean and standard deviation of digestive biscuits.

- 3 The times, in seconds, taken by 20 people to solve a simple numerical puzzle were

17 19 22 26 28 31 34 36 38 39
41 42 43 47 50 51 53 55 57 58

- (a) Calculate the mean and the standard deviation of these times.
- (b) In fact, 23 people solved the puzzle. However, 3 of them failed to solve it within the allotted time of 60 seconds.
Calculate the median and the interquartile range of the times taken by all 23 people.
- (c) For the times taken by all 23 people, explain why:
- (i) the mode is **not** an appropriate numerical measure.
 - (ii) the range is **not** an appropriate numerical measure.

Note: a question in such a context may ask students to find values of possible measures of central tendency and spread for a given set of data, and to critically assess which of these best represent the data.

L4

Recognise and interpret possible outliers in data sets and statistical diagrams.

Select or critique data presentation techniques in the context of a statistical problem.

Clean data, including dealing with missing data, errors and outliers.

Assessed at AS and A-level

Teaching guidance

Students should be able to:


- identify outliers either from a given rule or from observation of a given diagram.
- comment on the likely effect of removing the outlier.
- identify clear errors in data and comment on or suggest subsequent actions needed.
- select which of the representations in sections L1 and L2 is appropriate for representing given data sets.
- criticise, in context, a given representation.

Examples

- 1 Each Monday, Azher has a stall at a town's outdoor market. The table below shows, for each of a random sample of 10 Mondays during 2003, the air temperature, $x^{\circ}\text{C}$, at 9 am and Azher's takings, $\text{£}y$.

Monday	1	2	3	4	5	6	7	8	9	10
x	2	6	9	18	1	3	7	12	13	4
y	97	103	136	245	121	78	145	128	141	312

- (a) A scatter diagram of these data is shown below.



Give **two** distinct comments, in context, on what this diagram reveals.

- (b) One of the Mondays is found to be Easter Monday, the busiest Monday market of the year. Identify which Monday this is likely to be.

Note: in this question, students are left to decide what the salient features are that are being revealed on the scatter diagram. An ideal response would mention that there is a possible outlier (or that there are two possible outliers).

- 2 The table below shows data relating to the marital status of the adult populations of England and Wales during the period 1971 to 2006.

Mid-year	Total population	Males					Females				
		Single	Married	Divorced	Widowed	Total	Single	Married	Divorced	Widowed	Total
1971	36818	4173	12522	187	682	17563	3583	12566	296	2810	19255
1976	37486	4369	12511	376	686	17941	3597	12538		2877	19545
1981	38724	5013	12238	611	698	18559	4114	12284	828	2939	20165
1986	39837	5625	11867	917	695	19103	4617	12000	1165	2953	20734
1991	40501	5891	11636	1187	727	19441	4817	11833	1459	2951	21060
1996	40827	6225	11310	1346	733	19614	5168	11433	1730	2881	21212
2001	41865	6894	11090	1482	733	20198	5798	11150	1975	2745	21667
2006	43494	7833	10881	1696	716	21126	6683	10893	2244	2548	22367

Source: *Population trends*, office for National Statistics, 2009


- (a) (i) How many single males were there in England and Wales in 1981?
- (ii) The number of divorced females for 1976 has been omitted. Calculate the number which should be inserted in that space.
- (iii) The total population in the table for 1996 is **not** the sum of the total number of males and the total number of females.
- Assuming that the figures are correct, explain how this has happened.

Note: this question has used data that has been extracted from a much larger data set. This illustrates a possible way in which a large data set may be used.

- 3 The following table gives information on the average intake, per person per day, of Vitamin D, μg and Vitamin E, mg, for nine regions in England.

Region	Vitamin D	Vitamin E
North East	2.87	11.63
North West	3.09	11.92
Yorks and Humber	2.97	11.59
East Midlands	3.24	12.95
West Midlands	3.07	12.24
East	3.33	12.72
London	2.98	13.12
South East	3.16	12.09
South West	3.26	12.32

The scatter diagram illustrates this data.



- (a) Give two distinct comments on what the scatter diagram reveals.
- (b) The data for the London region is removed from the table and the data point for London is removed from the scatter diagram.

State what effect this would have on the correlation between average intake of Vitamin D and Vitamin E. Circle the correct answer.

Correlation would be weaker and positive

Correlation would stay the same

Correlation would be negative

Correlation would be stronger and positive

M

Probability

M1

Understand and use mutually exclusive and independent events when calculating probabilities.
Link to discrete and continuous distributions.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- find the probability of an event by extracting relevant information from a description of a situation (in context) or from a table of information.
- recognise and use set theory notation in the context of probability, eg $P(A \cup B)$, $P(A \cap B)$, $P(A')$
- recognise and define the meaning of mutually exclusive events, ie $P(A \cap B) = 0$
- define the condition for two events to be independent and determine whether two events are independent by finding, and comparing, relevant probabilities, ie $P(A \cap B) = P(A) \times P(B)$ or $P(A) = P(A|B)$, when the events A and B are independent.

Examples

- 1 Xavier, Yuri and Zara attend a sports centre for their judo club's practice sessions. The probabilities of them arriving late are, independently, 0.3, 0.4 and 0.2 respectively.
 - (a) Calculate the probability that for a particular practice session:
 - (i) all three arrive late.
 - (ii) none of the three arrives late.
 - (iii) only Zara arrives late.
 - (b) Zara's friend Wei also attends the club's practice sessions. The probability that Wei arrives late is 0.9 when Zara arrives late, and is 0.25 when Zara does not arrive late. Calculate the probability that for a particular practice session:
 - (i) both Zara and Wei arrive late.
 - (ii) either Zara or Wei, but not both, arrives late.

- 2 A school employs 75 teachers. The following table summarises their length of service at the school, classified by gender.

	Less than 3 years	3 years to 8 years	More than 8 years
Female	12	20	13
Male	8	15	7

- (a) Find the probability that a randomly selected teacher:
- (i) is female.
 - (ii) is female, given that the teacher has more than 8 years' service.
 - (iii) is female, given that the teacher has less than 3 years' service.
- (b) State, giving a reason, whether or not the event of selecting a female teacher is independent of the event of selecting a teacher with less than 3 years' service.
- (c) Define an event which is mutually exclusive to the event of selecting a female teacher.
- (d) Three teachers are selected at random without replacement, find the probability that all three are:
- (i) females with less than three years' service.
 - (ii) of the same gender.

- 3 A housing estate consists of 320 houses; 120 detached and 200 semi-detached. The numbers of children living in these houses are shown in the table.

	Number of children				
	None	One	Two	At least three	Total
Detached house	24	32	41	23	120
Semi-detached house	40	37	88	35	200
Total	64	69	129	58	320

A house on the state is selected at random.

D denotes the event 'the house is detached'.

R denotes the event 'no children live in the house'.

S denotes the event 'one child lives in the house'.

T denotes the event 'two children live in the house'.

(D' denotes the event 'not D ').

(a) Find:

(i) $P(D)$

(ii) $P(D \cap R)$

(b) (i) Name two of the events D , R , S and T that are mutually exclusive.

(ii) Determine whether the events D and R are independent.
Justify your answer.

(c) Define, in the context of this question, the event:

(i) $D' \cup T$

(ii) $D \cap (R \cup S)$

M2

Understand and use conditional probability, including the use of tree diagrams, Venn diagrams, two-way tables.

Understand and use the conditional probability formula

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Only assessed at A-level

Teaching guidance

Students should be able to:

- recognise that $P(A|B)$ is the probability that event A will happen, given that event B has already happened,
- find the probability $P(A|B)$ either by inspection or by using the probability formula $P(A \cap B) = P(A) \times P(B|A) = P(B) \times P(A|B)$ in a given context, which may include the use of tree diagrams, Venn diagrams and two-way tables.

Examples

- The British and Irish Lions 2005 rugby squad contained 50 players. The nationalities and playing positions of these players are shown in the table.

		Nationality			
		English	Welsh	Scottish	Irish
Playing position	Forward	14	5	2	6
	Back	8	7	2	6

A player was selected at random from the squad. Calculate the probability that the player was:

- English.
- Irish, given that the player was a back.
- a forward, given that the player was not Scottish.

- 2 The age and the blood pressure status for each adult male in a randomly selected sample are summarised in the following table.

Blood pressure status	Age range		
	25-34	35-54	55-74
Normal (untreated)	52	47	28
High (untreated)	4	7	18
High (treated)	1	4	13

One of the adult males is selected at random.

H is the event 'the male has high blood pressure'.

R is the event 'the male selected is aged 55-74'.

S is the event 'the male selected is aged 25-34'.

T is the event 'the male selected is being treated'.

Find:

- (a) $P(S)$
- (b) $P(T)$
- (c) $P(R \cup T)$
- (d) $P(H \cap S)$
- (e) $P(S | T)$
- (f) $P(R | H')$

- 3 Fred and his daughter, Delia, support their town's rugby team. The probability that Fred watches a game is 0.8. the probability that Delia watches a game is 0.9 when her father watches the game, and is 0.4 when her father does not watch the game.

(a) Calculate the probability that:

- (i) both Fred and Delia watch a particular game.
- (ii) neither Fred nor Delia watch a particular game.

- (b) Molly supports the same rugby team as Fred and Delia. The probability that Molly watches a game is 0.7, and is independent of whether or not Fred or Delia watches the game.

Calculate the probability that:

- (i) all three supporters watch a particular game.
- (ii) exactly two of the three supporters watch a particular game.

- 4 On a particular day, a sample of shoppers at a large supermarket were asked which, if any, of three categories of food, meat (M), dairy (D) and fruit and vegetables (F) they intended to buy on their visit that day.

Of these shoppers:

24 answered ' M '

60 answered ' D '

55 answered ' F '

45 answered ' M and D '

12 answered ' D and F '

8 answered ' M , D and F '

and 6 answered 'none of the categories'.

- (a) Draw a fully labelled Venn diagram to illustrate this information.
- (b) State the number of these shoppers intending to buy only one of the three categories of food.
- (c) Find $P(M \cap D | F')$

M3

Modelling with probability, including critiquing assumptions made and the likely effect of more realistic assumptions.

Only assessed at A-level

Teaching guidance

Students should be able to:

- assess and determine whether a stated probability model is appropriate in a given context.
- consider whether or not assumptions being made in order to use a given probability model are likely to be valid and the likely effect on results when more realistic assumptions are made.

Examples

- 1 A gas supplier maintains a team of engineers who are available to deal with leaks reported by customers. Most reported leaks can be dealt with fairly quickly but some require a long time. The time (excluding travelling time), X , taken to deal with reported leaks is found to have a mean of 65 minutes and a standard deviation of 60 minutes.

A statistician consulted by the gas supplier stated that, as the times had a mean of 65 minutes and a standard deviation of 60 minutes, the normal distribution would not provide an adequate model.

Explain the reason for the statistician's statement.

- 2 During June 2011, the volume, X litres, of unleaded petrol purchased per visit at a supermarket's filling station by private car customers could be modelled by a normal distribution with a mean of 32 and a standard deviation of 10.

(a) Determine:

(i) $P(X < 40)$

(ii) $P(X > 25)$

(iii) $P(25 < X < 40)$

(b) Given that during June 2011 unleaded petrol cost £1.34 per litre, calculate the probability that the unleaded petrol bill for a visit during June 2011 by a private car customer exceeded £65.

(c) Give **two** reasons, in any context, why the model $N(32, 10^2)$ is unlikely to be valid for a visit by **any** customer purchasing fuel at this filling station during June 2011.

- 3 The proportion of passengers who use senior citizen bus passes to travel into a particular town on park and ride buses between 9.30 am and 11.30 am on weekdays is 0.45.

It is proposed that, when there are n passengers on a bus, a suitable model for the number of passengers using senior citizen bus passes is the distribution $B(n, 0.45)$.

- (a) Assuming that this model applies to the 10.30 am weekday 'Park and Ride' bus into the town:
- (i) calculate the probability that, when there are **16** passengers, exactly three of them are using senior citizen bus passes.
 - (ii) determine the probability that, when there are **25** passengers, fewer than 10 of them are using senior citizen bus passes.
 - (iii) determine the probability that, when there are **40** passengers, at least 15 but at most 20 of them are using senior citizen bus passes.
 - (iv) calculate the mean and the variance for the number of passengers using senior citizen bus passes when there are **50** passengers.
- (b) (i) Give a reason why the proposed model may not be suitable.
- (ii) Give a **different** reason why the proposed model would not be suitable for the number of passengers using senior citizen bus passes to travel into the town on the **7.15 am** weekday Park and Ride bus.

N

Statistical distributions

N1

Understand and use simple, discrete probability distributions (calculation of mean and variance of discrete random variables is excluded), including the binomial distribution, as a model; calculate probabilities using the binomial distribution.

Assessed at AS and A-level

Teaching guidance

Students should:

- recognise when a situation may be modelled by a discrete random variable.
- know and be able to use the fact that the sum of the probabilities of all possible outcomes of an event is 1.
- be able to find the probability of a defined event in a given context.
- recognise and be able to use $B(n, p)$ as the notation for a binomial distribution with n independent trials where p is the probability of 'success' at any trial.
- be able to state the conditions necessary for a binomial distribution and assess whether they are likely to be valid in a given situation.
- be able to find the probability of an exact number of successes in a binomial distribution.
- be able to find cumulative probabilities in a binomial distribution.
- be able to use $P(\text{at least 1 'success'}) = 1 - P(\text{zero 'successes'})$.

Note: when using the binomial distribution, students will be expected to be able to use their calculator effectively to find directly both the probability of an exact number of outcomes and to find cumulative probabilities.

Examples

- 1 Todd is a dentist. Clients at Todd's surgery pay one of three possible fees: £20 for a check-up only, £50 for a check-up followed by minor treatment, and £210 for a check-up followed by major treatment. Experience shows that the probabilities for those needing treatment are as in the table.

	Fee	Probability
Check-up only	£20	
Check-up + minor treatment	£50	0.32
Check-up + major treatment	£210	0.11

- (a) Write down the probability for clients needing a check-up only.
- (b) Find the probability that a randomly selected patient in Todd's surgery will pay a fee that is less than £100.
- 2 An amateur tennis club purchases tennis balls that have been used previously in professional tournaments.
- The probability that each such ball fails a standard bounce test is 0.15.
- The club purchases boxes each containing 10 of these tennis balls. Assume that the 10 balls in any box represent a random sample.
- (a) Determine the probability that the number of balls in a box which fail the bounce test is:
- (i) at most 2.
 - (ii) at least 2.
 - (iii) more than 1 but fewer than 5.
- (b) Determine the probability that, in 5 boxes, the total number of balls which fail the bounce test is:
- (i) more than 5.
 - (ii) at least 5 but at most 10.

- 3 Each evening Aaron sets his alarm for 7 am. He believes that the probability that he wakes before his alarm rings each morning is 0.4, and is independent from morning to morning.

One week has seven mornings.

- (a) Assuming that Aaron's belief is correct, calculate values for the mean and standard deviation of the number of mornings in a week when Aaron wakes before his alarm rings.
- (b) During a 50-week period, Aaron records, each week, the number of mornings on which he wakes before his alarm rings. The results are as follows.

Number of mornings	0	1	2	3	4	5	6	7
Frequency	10	8	7	7	5	5	4	4

- (i) Calculate the mean and standard deviation of these data.
- (ii) State, giving reasons, whether your answers to part (b) (i) support Aaron's belief that the probability that he wakes before his alarm rings each morning is 0.4, and is independent from morning to morning.

Note: (b) (ii) assesses content from section M3 in that it requires students to critique the assumption that $p = 0.4$ and that it is independent from morning to morning.

N2

Understand and use the Normal distribution as a model; find probabilities using the Normal distribution.

Link to histograms, mean, standard deviation, points of inflection and the binomial distribution.

Only assessed at A-level

Teaching guidance

Students should:

- know that the normal distribution is a possible model for a continuous random variable.
- recognise and be able to use the notation $N(\mu, \sigma^2)$ to denote the normal distribution with population mean μ and population variance σ^2 .
- Know and be able to use the fact that areas under a normal distribution curve correspond to probabilities.
- know and be able to use the symmetry of the normal distribution (this may include using the knowledge that the population mean lies at the centre of a normal distribution).
- know and be able to use the property that the central 99.8% of a normal distribution lies within approximately three standard deviations either side of the mean of the distribution, and other similar results.
- understand that a z -score is a measure of how many standard deviations (σ) a value is to the right of the population mean and use the formula $z = \frac{x - \mu}{\sigma}$ to find a z -score.
- be able to find probabilities using the normal distribution.
- recognise when the symmetrical shape of a histogram would allow the normal distribution to be used as a model for the distribution and hence deduce approximate values of the mean and standard deviation of the population.
- recognise when the symmetry of a binomial distribution may permit the use of a normal distribution as a model and hence deduce approximate values of the mean and standard deviation of the population.
- know and be able to use and show that the points of inflection of a normal distribution lie one standard deviation either side of the mean of the distribution.
- recognise when a probability found using the normal distribution may be used as the value of p in a binomial distribution and subsequent use of binomial probabilities.

Note: students will be expected to find the values of probabilities in a normal distribution directly from their calculator.

Examples

- 1 The volume of Everwhite toothpaste in a pump-action dispenser may be modelled by a normal distribution with a mean of 106 ml and a standard deviation of 2.5 ml.
- Determine the probability that the volume of Everwhite in a randomly selected dispenser is:
- (a) less than 110 ml.
 - (b) more than 100 ml.
 - (c) between 104 ml and 108 ml.
 - (d) not exactly 106 ml.
- 2 Draught excluder for doors and windows is sold in rolls of nominal length 10 metres.
- The actual length, X metres, of draught excluder on a roll may be modelled by a normal distribution with mean 10.2 and standard deviation 0.15 such that $X \sim N(10.2, 0.15^2)$
- (a) Determine:
 - (i) $P(X < 10.5)$;
 - (ii) $P(10.0 < X < 10.5)$.
 - (b) A customer randomly selects six 10 metre rolls of the draught excluder.
- Calculate the probability that all six rolls selected contain more than 10 metres of draught excluder.

N3

Select an appropriate probability distribution for a context, with appropriate reasoning, including recognising when the binomial or Normal model may not be appropriate.

Only assessed at A-level

Teaching guidance

Students should:

- know and recognise that the binomial distribution may be used to model a discrete random variable.
- know, and be able to state, necessary and sufficient conditions for a discrete random variable to follow a binomial distribution and to recognise when one or more of these conditions may be being violated hence implying the binomial distribution is not a valid model.
- know and be able to use the mean of a binomial distribution is np and that the variance is $np(1 - p)$
- know and recognise that the normal distribution may be used to model a continuous random variable.
- know and be able to use the properties of the normal distribution, as given in section N2, and recognise when these are being violated hence implying the normal distribution is not a valid model.

Examples

- 1 In each situation below find the probability of the given event and state whether you are using the binomial distribution, the normal distribution or neither of these in each case.
 - (a) Past experience suggests that the weight of the contents jars of jam filled by a machine has a mean of 460.2 g and a standard deviation of 2.4 g.
Find the probability that the weight of the contents of a randomly selected jar of jam, filled by this machine, is at least 454.0 g and at most 464.0 g.
 - (b) A sports club has 50 members. Of these members 30 are adults. 15 are juniors and 5 are social members.
Three members are selected at random.
Find the probability that all three are junior members.
 - (c) A research paper suggests that 10% of the population of the UK is left-handed.
Find the probability that a random sample of 20 people, chosen from this population, contains at least 1 and at most 4 people who are left-handed.

-
- 2 A hotel has 50 single rooms, 16 of which are on the ground floor. The hotel offers guests a choice of a full English breakfast, a continental breakfast or no breakfast. The probabilities of these choices being made are 0.45, 0.25 and 0.30 respectively. It may be assumed that the choice of breakfast is independent from guest to guest.
- (a) On a particular morning there are 16 guests, each occupying a single room on the ground floor. Calculate the probability that exactly 5 of these guests require a full English breakfast.
 - (b) On a particular morning when there are 50 guests, each occupying a single room, determine the probability that:
 - (i) at most 12 of these guests require a continental breakfast;
 - (ii) more than 10 but fewer than 20 of these guests require no breakfast.
 - (c) When there are 40 guests, each occupying a single room, calculate the mean and the standard deviation for the number of guests requiring breakfast.

0

Statistical hypothesis testing

01

Understand and apply the language of statistical hypothesis testing, developed through a binomial model; null hypothesis; alternative hypothesis, significance level, test statistic, 1-tail test, 2-tail test, critical value, critical region, acceptance region, p -value; extend to correlation coefficients as measures of how close data points lie to a straight line and interpret a given correlation coefficient using a given p -value or critical value (calculation of correlation coefficients is excluded).

Assessed at AS and A-level

Teaching guidance

Students should:

- recognise whether a given context requires the use of a 1-tail or 2-tail test and understand the difference between them.
- be able to state appropriate null and alternative hypotheses to test a population proportion in a given context and know that the null hypothesis always contains the equality sign.
- understand that the significance level of a test is the probability of rejecting a correct null hypothesis in error.
- be able to find the test statistic as being the observed number of outcomes of the event.
- be able to find the critical region for a 1-tail test, or the critical regions for a 2-tail test, supporting the choice of values in such regions with appropriate binomial probabilities.
- know that the critical region consists of the critical values for the test and that if the test statistic lies in the critical region that this will lead to the rejection of the null hypothesis.
- know that the acceptance region is the range of possible values, that the discrete random variable can take, that do not lie in the critical region and that if the test statistic lies in the acceptance region that this will lead to the acceptance of the null hypothesis.
- appreciate that if the test statistic corresponds to a critical value in the critical region that the null hypothesis is rejected, or that if the test statistic is in the acceptance region then the null hypothesis is accepted.

- be able to use the given p -value corresponding to the test statistic or the given critical value(s), for the relevant significance level of the test, to decide whether to accept or reject the null hypothesis; understand that the p -value should be compared to a binomial distribution critical region with probability equal to or less than the significance level.
- be able to interpret a conclusion in context.

Examples

- 1 A test statistic has a binomial distribution, $B(25, 0.15)$.

Given that

$$H_0 : p = 0.15 \quad H_1 : p < 0.15$$

- (a) Find the critical region for the test statistic when the level of significance for the test is 5%.

Note: the critical region is the set of values of the test statistic that leads to the rejection of the null hypothesis, when the null hypothesis is assumed true. It is the area of the sampling distribution of a statistic that will lead to the rejection of the hypothesis tested when that hypothesis is true.

- (b) Find the actual probability of rejecting the null hypothesis when the critical region found in part (a) is used.

- 2 Previous experience suggests that 15% of the workers in a factory wear glasses.

Arwen selects a random sample of 20 workers from the factory and finds that 6 of them are wearing glasses. She suspects that the proportion of workers in the factory wearing glasses may have changed.

She uses the hypotheses

$$H_0 : p = 0.15 \quad H_1 : p > 0.15$$

and decides to use a 5% significance level for her test.

Her test statistic of '6 workers are wearing glasses' gives her a p -value of 0.06731.

Complete Arwen's test.

02

Conduct a statistical hypothesis test for the proportion in the binomial distribution and interpret the results in context.

Understand that a sample is being used to make an inference about the population and appreciate that the significance level is the probability of incorrectly rejecting the null hypothesis.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- state appropriate null and alternative hypotheses to test the value of a population proportion in a given context.
- find the test statistic as being the observed number of outcomes of the event.
- either find the critical region(s) for the test (supporting the choice of critical region with appropriate binomial probabilities) or find the p -value corresponding to the test statistic; understand that the p -value should be compared to a binomial critical region with probability equal to or less than the significance level.
- either compare the test statistic with the critical region or compare the p -value with the significance level of the test.
- state the conclusion in context.
- appreciate that the results obtained from the sample are being used to make an inference as to what is happening in the parent population and that the conclusion reached may be incorrect.

Examples

- 1 National records show that 35% of train passengers buy their tickets in advance. A random sample of 25 passengers using a particular railway station is selected, and it is found that 13 of them bought their tickets in advance.

Investigate, at the 10% level of significance, whether the data support the view that the percentage of passengers from this station who buy their tickets in advance is different from the national figure of 35%.

Note: the actual level of significance of this data is less than 10%.

- 2 James is a guitarist in a rock band which is about to start a 14-night tour. James usually uses Britepick guitar strings, which he changes before each performance. The thinnest string on a guitar, the top-E string, is the one most likely to break and, for James, the probability that this happens during a 1-hour performance is 0.02.
- (a) James is thinking of using Pluckwell strings rather than Britepick strings in the future and has bought some Pluckwell top-E strings to use each night of the 14-night tour. He finds that he breaks a top-E string during the band's 1-hour performance on two of these 14 nights.
- Investigate, at the 5% level of significance, whether Pluckwell top-E strings are more likely to break than Britepick top-E strings.
- (b) The band's manager, Noddy, suggests that the conclusion that James reached in part (a) is always going to be the same for these two makes of string.
- Comment on the validity of Noddy's suggestion.

Note: in part (b) students need to decide how best to approach this part of the question. Solutions may include that the conclusion is only based on the result arrived at by use of a small sample of strings and that this may not be true for all strings produced by the company – the breaking of strings may be influenced by temperature/humidity/striking the strings harder (or softer) on the tour. They may also include that with modern quality control the strings produced are likely to be extremely consistent and that although it is likely similar results will be achieved with similar testing conditions we can't say that the result will always be the same since there will be variability within output.

03

Conduct a statistical hypothesis test for the mean of a Normal distribution with known, given or assumed variance and interpret the results in context.

Only assessed at A-level

Teaching guidance

Students should be able to:

- state appropriate null and alternative hypotheses to test the value of a population mean in a given context.
- find the test statistic using the formula $\frac{\bar{X} - \mu}{\left(\frac{\sigma}{\sqrt{n}}\right)}$
- either find the critical value(s) for the test or find the p -value corresponding to the test statistic.
- either compare the test statistic with the critical value(s) or compare the p -value with the significance level of the test.
- decide whether to accept or reject the null hypothesis.
- state the conclusion in the context of the question.
- appreciate that the results obtained from the sample are being used to make an inference as to what is happening in the parent population and that the conclusion reached may be incorrect.

Examples

- As a special promotion, a supermarket offers cartons of orange juice containing '25% extra' with no price increase.

A random sample of cartons of orange juice was checked. The percentages by which the contents exceeded the nominal quantity were recorded, with the following results:

23.3 27.5 25.7 20.9 24.3 22.6 21.5 22.1

Examine whether the mean percentage by which the contents exceed the nominal quantity is less than 25. Use the 5% significance level. Assume that the data are from a normal distribution with standard deviation 2.3.

- A machine fills paper bags with flour. Before maintenance on the machine, the weight of the flour in a bag could be modelled by a normal distribution with mean 1005 g and standard deviation 2.1 g. Following this maintenance, the flour in each of a random sample of 8 bags was weighed. The weights, in grams, were as follows:

1006.1 1004.9 1005.8 1007.9 1004.7 1006.3 1007.4 1007.2

Carry out a test at the 10% significance level, to decide whether the mean weight of flour in a bag filled by the machine had **changed**. Assume that the distribution of weights was still normal with standard deviation 2.1 g.

P

Quantities and units in mechanics

P1

Understand and use fundamental quantities and units in the SI system: length, time, mass.

Understand and use derived quantities and units: velocity, acceleration, force, weight and moment.

Assessed at AS and A-level

Teaching guidance

Students should:

- know and be able to use the following:

Fundamental quantity	SI base unit
length	metre (m)
time	second (s)
mass	kilogram (kg)

- be able to convert between commonly used SI units, for example kilometres and metres or kilograms and tonnes.
- know and be able to use the following:

Derived quantity	SI unit
velocity	metre per second (m s^{-1})
acceleration	metre per second squared (m s^{-2})
force/weight	newton (N)
moment	newton metre (Nm) [A-level only]

- understand that g is acceleration due to gravity (m s^{-2}).
- understand that weight is a force, $W = mg$ (N).
- if required, be able to convert non-standard units, for example kilometres per hour, to metres per second.
- be familiar with equivalent notions such as m/s^2 and m s^{-2}
- know that trigonometric ratios and the coefficient of friction are dimensionless.

Example

- 1 A car of mass 1.2 tonnes is accelerating at 5.0 m s^{-2} .


Calculate the resultant force acting on the car.

Only assessed at A-level

Example

- 1 A uniform rod, of length 1.20 metres, has a mass of 3.00 kg. The rod is held in equilibrium by two supports, A and B, as shown in the diagram.

Calculate the clockwise moment of the weight of the rod about support A.



Circle your answer.

45 Nm

3.6 kg/m

1.35g Nm

3g N

Note: in mechanics questions where measured quantities have been given to varying degrees of accuracy students should give their final answer to an appropriate degree of accuracy. For example, if some measured lengths are given to two significant figures, but other measured lengths are given to three significant figures the final answer should be given to two significant figures.

Q

Kinematics

Q1

Understand and use the language of kinematics: position; displacement; distance travelled; velocity; speed; acceleration.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand positions described relative to a given origin.
- understand and describe the position of a particle through a combination of its initial position and a displacement.
- demonstrate an understanding of the relationship between the vectors displacement and velocity and their associated scalar quantities distance and speed.
- understand average speed and average velocity.

Examples

- 1 A car travels on a straight horizontal race track. The car decelerates uniformly from a speed of 20 m s^{-1} to a speed of 12 m s^{-1} as it travels a distance of 640 metres. The car then accelerates uniformly, travelling a further 1820 metres in 70 seconds.
- (a) (i) Find the time that it takes the car to travel the first 640 metres.
- (ii) Find the deceleration of the car during the first 640 metres.
- (b) (i) Find the acceleration of the car as it travels the further 1820 metres.
- (ii) Find the speed of the car when it has completed the further 1820 metres.

- 2 Two boys, Alf and Bob, are running a 100m race. After 4 seconds Alf has run 21.5 metres and Bob has run 18.3 metres and both boys have reached their maximum speeds. The boys complete the race running at their maximum speeds. Alf's maximum speed is 5.9 m s^{-1} and Bob's maximum speed is 6.2 m s^{-1} .

Determine:

- (a) the total time taken for the race to be won.
- (b) the distance from the finish of the boy in second place when the race is won.

Note: students should be encouraged to give final answers to an appropriate degree of accuracy.

Q2

Understand, use and interpret graphs in kinematics for motion in a straight line: displacement against time and interpretation of gradient; velocity against time and interpretation of gradient and area under the graph.


Assessed at AS and A-level

Teaching guidance

Students should be able to:

- use the gradient of a displacement-time graph to find the velocity (or speed).
- use the gradient of a velocity-time graph to give the acceleration and interpret positive and negative gradients.
- understand that graphs may include negative velocities.
- use the area under a velocity-time graph to find displacements.
- sketch either a displacement-time or velocity-time graph for a given scenario.

- 1 The diagram shows a velocity-time graph for a train as it moves on a straight horizontal track for 50 seconds.



- Find the distance that the train moves in the first 28 seconds.
- Calculate the total distance moved by the train during the 50 seconds.
- Find the displacement of the train from its initial position when it has been moving for 50 seconds.
- Find the acceleration of the train in the first 18 seconds of its motion.

2 A van moves from rest on a straight horizontal road.


- (a) In a simple model, the first 30 seconds of the motion are represented by three separate stages, each lasting 10 seconds and each with a constant acceleration.

During the first stage, the van accelerates from rest to a velocity of 4 m s^{-1} .

During the second stage, the van accelerates from 4 m s^{-1} to 12 m s^{-1} .

During the third stage, the van accelerates from 12 m s^{-1} to 16 m s^{-1} .


- (i) Sketch a velocity-time graph to represent the motion of the van during the first 30 seconds of its motion.
 - (ii) Find the total distance that the van travels during the 30 seconds.
 - (iii) Find the greatest acceleration of the van during the 30 seconds.
- (b) In another model of the 30 seconds of the motion, the acceleration of the van is assumed to vary during the first and third stages of the motion, but to be constant during the second stage, as shown in the velocity-time graph below.



The velocity of the van takes the same values at the beginning and the end of each stage of the motion as in part (a).

- (i) State, with a reason, whether the distance travelled by the van during the first 10 seconds of the motion in **this** model is greater or less than the distance travelled during the same interval in the model in part (a).
- (ii) Give one reason why **this** model represents the motion of the van more realistically than the model in part (a).

- 3 The graph shows the velocity of a parachutist t seconds after jumping from a hot-air balloon, before she opens her parachute.



What is the value of t when her acceleration is at a maximum?

Q3

Understand, use and derive the formulae for constant acceleration for motion in a straight line; extend to two dimensions using vectors.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- recall and use the following formulae:

$$v = u + at$$

$$s = \frac{1}{2}(u + v)t$$

$$s = ut + \frac{1}{2}at^2$$


$$s = vt - \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

Note: the less fashionable $s = vt - \frac{1}{2}at^2$ is not essential.

- derive the above formulae starting from given assumptions.

This may include starting from a graph similar to the one below.



Key

s = displacement
 u = initial velocity
 v = final velocity
 a = acceleration
 t = time

For example, the shaded area gives the displacement, s , and can be found as the area of a trapezium:

$$s = \frac{1}{2}(u + v)t$$

Example

- 1 Two cameras record the time that it takes a car on a motorway to travel a distance of 2 km. A car passes the first camera travelling at 32.0 ms^{-1} . The car continues at this speed for 12.5 seconds and then decelerates uniformly until it passes the second camera when its speed has decreased to 18.0 ms^{-1} .
- (a) Calculate the distance travelled by the car in the first 12.5 seconds.
 - (b) Find the time for which the car is decelerating.
 - (c) Sketch a speed-time graph for the car on this 2 km stretch of motorway.
 - (d) Find the total time taken for the car to travel along this 2 km stretch of motorway.

Only assessed at A-level

Teaching guidance

Students should be able to extend linear constant acceleration equations using the notation:

\mathbf{r} = position vector
 \mathbf{r}_0 = initial position
 \mathbf{u} = initial velocity
 \mathbf{v} = final velocity
 \mathbf{a} = acceleration
 t = time

Linear constant acceleration equation	Vector constant acceleration equation
$v = u + at$	$\mathbf{v} = \mathbf{u} + \mathbf{a}t$
$s = \frac{1}{2}(u + v)t$	$\mathbf{r} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t + \mathbf{r}_0$
$s = ut + \frac{1}{2}at^2$	$\mathbf{r} = \mathbf{u}t + \frac{1}{2}\mathbf{a}t^2 + \mathbf{r}_0$
$s = vt - \frac{1}{2}at^2$	$\mathbf{r} = \mathbf{v}t - \frac{1}{2}\mathbf{a}t^2 + \mathbf{r}_0$
$v^2 = u^2 + 2as$	No equivalent

Note: when working with vectors, students can use column vectors $\begin{pmatrix} a \\ b \end{pmatrix}$ or express them in the form $a\mathbf{i} + b\mathbf{j}$.

Example

- 1 A helicopter is initially hovering above a lighthouse. It then sets off so that its acceleration is $(0.500\mathbf{i} + 0.375\mathbf{j}) \text{ ms}^{-2}$. The helicopter does not change its height above sea level as it moves. The unit vectors \mathbf{i} and \mathbf{j} are directed east and north respectively.
- (a) Find the speed of the helicopter 20.0 seconds after it leaves its position above the lighthouse.
 - (b) Find the bearing on which the helicopter is travelling, giving your answer correct to the nearest degree.
 - (c) The helicopter stops accelerating when it is 500 metres from its initial position. Find the time that it takes for the helicopter to travel from its initial position to the point where it stops accelerating.

Q4

Use calculus in kinematics for motion in a straight line:

$v = \frac{dr}{dt}$, $a = \frac{dv}{dt} = \frac{d^2r}{dt^2}$, $r = \int v dt$, $v = \int a dt$; extend to two dimensions using vectors.

Assessed at AS and A-level

Teaching guidance

Students should:

- know and be able to apply the following to motion in a straight line:

$$v = \frac{dr}{dt} \qquad a = \frac{dv}{dt} = \frac{d^2r}{dt^2}$$

$$r = \int v dt \qquad v = \int a dt$$

- explore the relationship between calculus methods and the constant acceleration equations.

Note: at AS, questions will be limited to integrating or differentiating functions formed from sums and differences of terms such as At^n (with $n \neq -1$ in the case of integration), matching the level of difficulty required for sections G and H at AS.

Examples

- A particle moves in a straight line and at a time t it has velocity v , where

$$v = 3t^2 - 12t + 6$$

- (i) Find an expression for the acceleration of the particle at time t .

- (ii) When $t = 2$, show that the acceleration of the particle is 0.

- When $t = 0$, the particle is at the origin.

Find an expression for the displacement of the particle from the origin at time t .

- A particle moves in a straight line with constant acceleration, a .

Given that the initial velocity of the particle is u , use integration to prove that the displacement of the particle from its initial position is given by

$$s = ut + \frac{1}{2}at^2$$

Only assessed at A-level

Teaching guidance

Students should be able to:

- answer questions that draw on any of the differentiation or integration techniques from the pure mathematics sections G and H, for example:
 - find the velocity, given that the acceleration, a , at time t is given by $a = t^2 e^{-t}$
 - find the acceleration given that the displacement, s , at a time t is given by $s = e^{\frac{-t}{2}} \sin\left(\frac{nt}{20}\right)$
- recall and use the facts that in two dimensions, if $\mathbf{r} = x\mathbf{i} + y\mathbf{j}$ then:

$$\mathbf{v} = \frac{dx}{dt}\mathbf{i} + \frac{dy}{dt}\mathbf{j} \text{ and } \mathbf{a} = \frac{d^2x}{dt^2}\mathbf{i} + \frac{d^2y}{dt^2}\mathbf{j}$$

Examples

- 1 A particle moves in a straight line and at time t it has velocity v , where

$$v = 3t^2 - 2 \sin 3t + 6$$

- (a) (i) Find an expression for the acceleration of the particle at time t .

- (ii) When $t = \frac{\pi}{3}$, show that the acceleration of the particle is $2\pi + 6$.

- (b) When $t = 0$, the particle is at the origin.

Find an expression for the displacement of the particle from the origin at time t .

- 2 A particle moves on a horizontal plane, in which the unit vectors \mathbf{i} and \mathbf{j} are directed east and north respectively.

At time t seconds, the position vector of the particle is \mathbf{r} metres, where

$$\mathbf{r} = \left(2e^{\frac{1}{2}t} - 8t + 5\right)\mathbf{i} + (t^2 - 6t)\mathbf{j}$$

- (a) Find an expression for the velocity of the particle at time t .

- (b) (i) Find the speed of the particle when $t = 3$.

- (ii) State the direction in which the particle is travelling when $t = 3$.

- (c) Find the acceleration of the particle when $t = 3$.

- (d) The mass of the particle is 7 kg. Find the magnitude of the resultant force on the particle when $t = 3$.

Q5

Model motion under gravity in a vertical plane using vectors; projectiles.

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- use the acceleration due to gravity in the vector form:

$$\mathbf{a} = -g \mathbf{j} \quad \text{or} \quad \mathbf{a} = \begin{pmatrix} 0 \\ -g \end{pmatrix}$$

where \mathbf{j} is defined as vertically upwards from the earth's surface.

Notes: questions could be set in the context of vertical motion or in two dimensions.

In projectile questions, students can expect a wide range of scenarios.

When solving equations:

- calculators may be used to solve quadratic equations. It is recommended that both solutions are always given by students and the appropriate solution selected, with explicit justification for the selection.
- students may need to use the following trigonometric identities:

$$\sin 2\theta = 2 \sin \theta \cos \theta$$

$$\sec^2 \theta = 1 + \tan^2 \theta$$

- use the value of 9.8, 9.81 or 10 for g , as directed in the question.
- use and understand assumptions made when modelling projectiles. For example:
 - projectile is a particle (has no size and does not spin).
 - projectile does not experience air resistance or wind.
 - projectile is launched and lands at the same level.


Note: in questions where a numerical value for g is needed, students will be clearly told which approximation to use and their answers should then be given to an appropriate degree of accuracy. When deciding on the degree of accuracy to use in their answers, students should be guided by the accuracy of the data given in the question.

Examples


- 1 In this question use $g = 9.8 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.

A ball is hit by a bat so that, when it leaves the bat, its velocity is 40 m s^{-1} at an angle of 35° above the horizontal. Assume that the ball is a particle and that its weight is the only force that acts on the ball after it has left the bat.

- (a) A simple model assumes that the ball is hit from the point A and lands for the first time at the point B , which is at the same level as A , as shown in the diagram.



- (i) Show that the time that it takes for the ball to travel from A to B is 4.7 seconds, correct to two significant figures.
- (ii) Find the horizontal distance from A to B .
- (b) A revised model assumes that the ball is hit from the point C , which is 1 metre above A . the ball lands at the point D , which is at the same level as A , as shown in the diagram.



Find the time that it takes for the ball to travel from C to D .

Note that in part (b) only one significant figure accuracy is appropriate.

- 2 In this question use $g = 10 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.

A projectile is launched at ground level, on a horizontal surface, with speed V at an angle θ above the horizontal.


- Show that the range of the projectile is $\frac{V^2 \sin 2\theta}{g}$ and state any essential assumptions that you make to obtain this result.
- Find an expression for the maximum height of the projectile.
- Calculate the range of a projectile launched at a speed of 30 m s^{-1} and at an angle of 40° above the horizontal.

- 3 In this question use $g = 10 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.

A ball is thrown at an angle α above the horizontal with an initial speed of $V \text{ m s}^{-1}$. At time t seconds the horizontal displacement of the ball from its initial position, O , is x metres and the vertical displacement is y metres. Assume that the only force acting on the ball after it has been thrown is its weight.

- Show that $y = x \tan \alpha - \frac{gx^2}{2V^2}(1 + \tan^2 \alpha)$

The ball is thrown from O , so that it passes through two small hoops. The hoops are set at horizontal distances of 2 m and 4 m from O . The positions of the hoops are shown in the diagram. Assume that the ball is a particle that passes through the centre of each hoop.



- Show that $\tan \alpha = \frac{13}{4}$ and find V
- If a heavier ball was thrown with the same initial velocity, would it pass through the hoops? Give a reason for your answer.

R

Forces and Newton's laws

R1

Understand the concept of a force; understand and use Newton's first law.

Assessed at AS and A-level

Teaching guidance

Students should:


- understand types of force, including:
 - normal reaction force
 - tension in a string or a rod
 - thrust in a rod
 - weight
 - friction.
- know that the resultant force acting on a body is zero if a body is in equilibrium.
- be able to find unknown forces acting on bodies that are at rest or moving with constant velocity.
- be able to model forces as vectors
- be able to find the resultant of several forces acting at a point.

Notes: students may be required to express a resultant force using components of a vector. Students may be required to find the magnitude and direction of a resultant force expressed as a vector.

Students will not be required to resolve forces at AS.

Examples

- 1 The diagram shows three forces and the perpendicular unit vectors \mathbf{i} and \mathbf{j} , which all lie in the same plane.



- Express the resultant of the three forces in terms of \mathbf{i} and \mathbf{j} .
 - Find the magnitude of the resultant force.
 - Draw a diagram to show the direction of the resultant force, and find the angle that it makes with the unit vector \mathbf{i} .
- 2 A car is travelling with a constant velocity of 15 m s^{-1} , in a straight line, on a horizontal road. A driving force of 600 N acts in the direction of motion and a resistance force opposes the motion of the car. Assume that no other horizontal forces act on the car.
- What is the magnitude of the resistance force acting on the car?
- Circle the correct answer.

600 N

40 N

9000 N

0 N

Only assessed at A-level

Teaching guidance


Students should be able to:

- model forces as vectors
- resolve forces, but only in two dimensions.

Example

- 1 In this question, use $g = 10 \text{ m s}^{-2}$, giving your final answer to an appropriate degree of accuracy.

A child pulls a sledge, of mass 8 kg , along a rough horizontal surface, using a light rope. The coefficient of friction between the sledge and the surface is 0.3 . The tension in the rope is T newtons. The rope is kept at an angle of 30° to the horizontal, as shown in the diagram.



Model the sledge as a particle.

- Draw a diagram to show all the forces acting on the sledge.
- Find the magnitude of the normal reaction force acting on the sledge in terms of T .
- Given that the sledge moves with a constant velocity, find the value of T .

R2

Understand and use Newton's second law for motion in a straight line (restricted to forces in two perpendicular directions or simple cases of forces given as 2-D vectors); extend to situations where forces need to be resolved (restricted to two dimensions).

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- use $F = ma$ for constant mass and constant force.
- understand that objects can be modelled as particles.
- comment on the relevance of any modelling assumptions made.

Note: questions may be set that require the use of constant acceleration equations together with Newton's second law.

Examples

- 1 A lift rises vertically from rest with constant acceleration.
After 4 seconds, it is moving upwards with a velocity of 2 m s^{-1} .
It then moves with a constant velocity for 5 seconds.
The lift then slows down uniformly, coming to rest after it has been moving for a total of 12 seconds.
 - (a) Sketch a velocity-time graph for the motion of the lift.
 - (b) Calculate the total distance travelled by the lift.
 - (c) The lift is raised by a single vertical cable. The mass of the lift is 300 kg.
Find the maximum tension in the cable during this motion.
- 2 Three forces act on a particle. These forces are $(9\mathbf{i} - 3\mathbf{j})$ newtons, $(5\mathbf{i} + 8\mathbf{j})$ newtons and $(-7\mathbf{i} + 3\mathbf{j})$ newtons. The vectors \mathbf{i} and \mathbf{j} are perpendicular unit vectors.
 - (a) Find the resultant of these forces.
 - (b) Find the magnitude of the resultant force.
 - (c) Given that the particle has mass 5 kg, find the magnitude of the acceleration of the particle.
 - (d) Find the angle between the resultant force and the unit vector \mathbf{i} .

Only assessed at A-level


Teaching guidance

Students should be able to answer questions set on an inclined plane or in other contexts that require forces to be resolved.

Examples

- 1 In this question, use $g = 9.8 \text{ m s}^{-2}$, giving your final answer to an appropriate degree of accuracy.

Three forces act in a vertical plane on an object of mass 250 g , as shown in the diagram.



The two forces P newtons and Q newtons each act at 80° to the horizontal. The object accelerates horizontally at $a \text{ m s}^{-2}$ under the action of these forces.

- (a) Show that

$$P = 125 \left(\frac{a}{\cos 80^\circ} + \frac{g}{\sin 80^\circ} \right)$$

- (b) Find the value of a for which Q is zero.

R3

Understand and use weight and motion in a straight line under gravity; gravitational acceleration, g , and its value in SI units to varying degrees of accuracy.

(The inverse square law for gravitation is not required and g may be assumed to be constant; students should be aware that g is not a universal constant but depends on location).

Assessed at AS and A-level

Teaching guidance

Students should be able to:

- understand the distinction between mass and weight.

Notes: In questions where a numerical value for g is needed, students will be clearly told which approximation to use and their answers should then be given to an appropriate degree of accuracy. When deciding on the degree of accuracy to use in their answers, students should be guided by the accuracy of the data given in the question.

In questions involving objects in motion under gravity it will be assumed that:

- g remains constant
- objects can be treated as particles
- resistance forces are negligible
- state necessary modelling assumptions, if required, that they have made and comment on how these assumptions relate to the model used.

Examples

- 1 In this question, use $g = 10 \text{ m s}^{-2}$, giving your final answer to an appropriate degree of accuracy.

A stone is dropped from a high bridge and falls vertically.

- (a) Find the distance that the stone falls during the first 4 seconds of its motion.
- (b) Find the speed of the stone after the first 4 seconds of its motion.
- (c) State one modelling assumption that you have made about the forces acting on the stone during the motion.

Note: in this question we would expect final answers given to one significant figure as all data given in the question is no more accurate than this.

- 2** In this question, use $g = 9.8 \text{ m s}^{-2}$, giving your final answer to an appropriate degree of accuracy.

On earth, an astronaut wearing her full spacesuit weighs 2100 newtons.

On the moon, wearing the same spacesuit, she weighs 350 newtons.

Calculate the acceleration due to gravity on the moon.

Note: in this question an answer with an accuracy of two significant figures would be appropriate.

R4

Understand and use Newton's third law; equilibrium of forces on a particle and motion in a straight line (restricted to forces in two perpendicular directions or simple cases of forces given as 2-D vectors); application to problems involving smooth pulleys and connected particles; resolving forces in two dimensions; equilibrium of a particle under coplanar forces.

Assessed at AS and A-level

Teaching guidance

Connected particles

Students should:

- understand that usually strings will be modelled as light and inextensible.
- understand that usually pulleys will be modelled as light and smooth.
- understand that usually pegs will be modelled as smooth.

Notes: Questions can be set involving objects that can be modelled as particles and are connected by a light, inextensible string.

Questions can be set that involve contexts such as a car towing a trailer or several carriages connected together as a train.

At AS, questions will be restricted to connected particles that move horizontally or vertically. Questions involving inclined planes will **not** be set.


Newton's third law

Students should be able to identify 'action and reaction' forces, including:

- weights
- normal reaction forces
- friction forces.

Examples


- 1 A wooden block, of mass 4 kg, is placed on a rough horizontal surface. The coefficient of friction between the block and the surface is 0.3. A horizontal force, of magnitude 30 newtons, acts on the block and causes it to accelerate.



Draw a diagram to show all the forces acting on the block.

- 2 In this question, use $g = 10 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.


Two particles, A and B, have masses 4 kg and 6 kg respectively. They are connected by a light inextensible string that passes over a smooth fixed peg. A second light inextensible string is attached to A. The other end of the string is attached to the ground directly below A. The system remains at rest, as shown in the diagram.



- (a) (i) Write down the tension in string connecting A and B.
- (ii) Find the tension in the string connecting A to the ground.
- (b) The string connecting particle A to the ground is cut. Find the acceleration of A after the string has been cut.


- 3 In this question, use $g = 9.8 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.

A block, of mass 14 kg , is held at rest on a rough horizontal surface. The coefficient of friction between the block and the surface is 0.25 . A light inextensible string, which passes over a fixed smooth peg, is attached to the block. The other end of the string is attached to a particle, of mass 6.0 kg , which is hanging at rest.



The block is released and begins to accelerate.

- Find the magnitude of the friction force acting on the block.
 - By forming two equations of motion, one for the block and one for the particle, show that the magnitude of the acceleration of the block and the particle is 1.2 m s^{-2} .
 - Find the tension in the string.
 - When the block is released, it is 0.80 metres from the peg. Find the speed of the block when it hits the peg.
 - When the block reaches the peg, the string breaks and the particle falls a further 0.50 metres to the ground. Find the speed of the particle when it hits the ground.
- 4 A small train at an amusement park consists of an engine and two carriages connected to each other by light horizontal rods, as shown in the diagram.



The engine has mass 2000 kg and each carriage has mass 500 kg .

The train moves along a straight horizontal track. A resistance force of magnitude 400 newtons acts on the engine, and resistance forces of magnitude 300 newtons act on each carriage. The train is accelerating at 0.50 m s^{-2} .

- Draw a diagram to show the **horizontal** forces acting on Carriage 2.
- Show that the magnitude of the force that the rod exerts on carriage 2 is 550 newtons.
- Find the magnitude of the force that rod attached to the engine exerts on Carriage 1.
- A forward driving force of magnitude P newtons acts on the engine. Find P .

Only assessed at A-level

Teaching guidance


Students should be able to:

- answer questions that involve resolving forces.
- understand that motion may not be restricted to horizontal or vertical and that inclined planes may be used.

Example

- 1 In this question, use $g = 9.81 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.

A particle, of mass 10.5 kg , is suspended in equilibrium by two light strings, AP and BP . The string AP makes an angle of 32.5° to the horizontal and the other string, BP , is horizontal, as shown in the diagram.



- Draw and label a diagram to show the forces acting on the particle.
- Show that the tension in the string AP is 192 N , correct to three significant figures.
- Find the tension in the horizontal string BP .

R5

Understand and use addition of forces; resultant forces; dynamics for motion in a plane.

Only assessed at A-level


Teaching guidance

Students should be able to:


- find resultants by use of a vector diagram or resolving into perpendicular components.
Note: unless specifically stated in the question, any appropriate method for finding the resultant is acceptable.
- use $F = ma$ in the form $F = m \frac{dv}{dt}$ to set up and solve a differential equation.
Note: mass will be constant.

Examples

- 1 Two forces, acting at a point, have magnitudes of 40 newtons and 70 newtons. The angle between the two forces is 30° , as shown in the diagram.



- (a) Find the magnitude of the resultant of these two forces.
- (b) Find the angle between the resultant force and the 70 newton force.
- 2 Three forces \mathbf{F}_1 , \mathbf{F}_2 and \mathbf{F}_3 act on a particle of mass 1.5 kg, as shown in the diagram.



Find the magnitude and direction of the acceleration of the particle.

3 In this question use $g = 9.8 \text{ m s}^{-2}$.

Vicky has mass 65 kg and is skydiving. She steps out of a helicopter and falls vertically. She then waits a short period of time before opening her parachute. The parachute opens at time $t = 0$ when her speed is 19.6 m s^{-1} . She then experiences an air resistance force of magnitude $260v$ newtons, where $v \text{ m s}^{-1}$ is her speed at time t seconds.

(a) When $t > 0$:

(i) Show that the resultant downward force acting on Vicky is $65(9.8 - 4v)$ newtons.

(ii) Show that $\frac{dv}{dt} = -4(v - 2.45)$

(b) By showing that

$$\int \frac{1}{v - 2.45} dv = -\int 4 dt$$

find v in terms of t .

R6

Understand and use the $\mathbf{F} \leq \mu \mathbf{R}$ model for friction; coefficient of friction; motion of a body on a rough surface; limiting friction and statics.

Only assessed at A-level

Teaching guidance

Students should be able to:

- answer questions asked in the context of inclined planes.
- understand that particles may be in equilibrium, limiting equilibrium or accelerating.
- find the range of possible values for μ .
- understand when $\mathbf{F} = \mu \mathbf{R}$ can be used.

- 1 In this question use $g = 9.8 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.


A sledge of mass 8.00 kg is at rest on a rough horizontal surface. A child tries to move the sledge by pushing it with a pole, as shown in the diagram, but the sledge **does not move**. The pole is at an angle of 30° to the horizontal and exerts a force of 40 newtons on the sledge.



- Draw a diagram to show the four forces acting on the sledge.
- Show that the normal reaction force between the sledge and the surface has magnitude 98 N .
- Find the magnitude of the friction force that acts on the sledge.
- Find the smallest possible value of the coefficient of friction between the sledge and the surface.

- 2 In this question use $g = 9.8 \text{ m s}^{-2}$, giving your final answers to an appropriate degree of accuracy.

A rough slope is inclined at an angle of 25° to the horizontal. A box of weight 80 newtons is on the slope. A rope is attached to the box and is parallel to the slope. The tension in the rope is of magnitude T newtons. The diagram shows the slope, the box and the rope.



- (a) The box is held in equilibrium by the rope.
- (i) Show that the normal reaction force between the box and the slope is 73 newtons.
 - (ii) The coefficient of friction between the box and the slope is 0.32. Find the magnitude of the maximum value of the frictional force which can act on the box.
 - (iii) Find the least possible tension in the rope to prevent the box from moving down the slope.
 - (iv) Find the greatest possible tension in the rope.
 - (v) Show that the mass of the box is approximately 8.2 kg.
- (b) The rope is now released and the box slides down the slope. Find the acceleration of the box.

S

Moments

S1


Understand and use moments in simple static contexts.

Only assessed at A-level

Teaching guidance

Students should:


- know that the centre of mass of uniform beams and rectangular laminae can be determined by symmetry.
- be able to answer questions in which forces act in perpendicular directions. For example, to determine the resultant moment of a situation similar to the one below:



Example

- 1 Ken is trying to cross a river of width 4.0 m. He has a uniform plank, AB , of length 8.0 m and mass 17 kg. The ground on both edges of the river bank is horizontal. The plank rests at two points, C and D , on fixed supports which are on opposite sides of the river. The plank is at right angles to both river banks and is horizontal. The distance AC is 1.0 m, and the point C is at a horizontal distance 0.60 m from the river bank.

Ken, who has mass 65 kg, stands on the plank directly above the middle of the river, as shown in the diagram.



- (a) Draw a diagram to show the forces acting on the plank.
- (b) Given that the reaction on the plank at the point D is $44g$ N, where g is the acceleration due to gravity, find the horizontal distance of the point D from the nearest bank.
- (c) State how you have used the fact that the plank is uniform in your solution.

A1

Appendix A

Mathematical notation for AS and A-level qualifications in Mathematics and Further Mathematics

The tables below set out the notation that must be used by AS and A-level Mathematics and Further Mathematics specifications. Students will be expected to understand this notation without need for further explanation.

Mathematics students will not be expected to understand notation that relates only to Further Mathematics content. Further Mathematics students will be expected to understand all notation in the tables.

For Further Mathematics, the notation for the core content is listed under sub-headings indicating 'Further Mathematics only'.

AS students will be expected to understand notation that relates to AS content, and will not be expected to understand notation that relates only to A-level content.

1	Set Notation	
1.1	\in	is an element of
1.2	\notin	is not an element of
1.3	\subseteq	is a subset of
1.4	\subset	is a proper subset of
1.5	$\{x_1, x_2, \dots\}$	the set with elements x_1, x_2, \dots
1.6	$\{x : \dots\}$	the set of all x such that ...
1.7	$n(A)$	the number of elements in set A
1.8	\emptyset	the empty set
1.9	\mathcal{E}	the universal set
1.10	A'	the complement of the set A
1.11	\mathbb{N}	the set of natural numbers, $\{1, 2, 3, \dots\}$

1.12	\mathbb{Z}	the set of integers, $\{0, \pm 1, \pm 2, \pm 3, \dots\}$
1.13	\mathbb{Z}^+	the set of positive integers, $\{1, 2, 3, \dots\}$
1.14	\mathbb{Z}_0^+	the set of non-negative integers, $\{0, 1, 2, 3, \dots\}$
1.15	\mathbb{R}	the set of real numbers
1.16	\mathbb{Q}	the set of rational numbers, $\left\{\frac{p}{q} : p \in \mathbb{Z}, q \in \mathbb{Z}^+\right\}$
1.17	\cup	union
1.18	\cap	intersection
1.19	(x, y)	the ordered pair x, y
1.20	$[a, b]$	the closed interval $\{x \in \mathbb{R} : a \leq x \leq b\}$
1.21	$[a, b)$	the interval $\{x \in \mathbb{R} : a \leq x < b\}$
1.22	$(a, b]$	the interval $\{x \in \mathbb{R} : a < x \leq b\}$
1.23	(a, b)	the open interval $\{x \in \mathbb{R} : a < x < b\}$

1	Set Notation (Further Mathematics only)	
1.24	\mathbb{C}	the set of complex numbers

2	Miscellaneous symbols	
2.1	$=$	is equal to
2.2	\neq	is not equal to
2.3	\equiv	is identical to or is congruent to
2.4	\approx	is approximately equal to
2.5	∞	infinity
2.6	\propto	is proportional to
2.7	\therefore	therefore

2.8	\because	because
2.9	$<$	is less than
2.10	\leq	is less than or equal to, is not greater than
2.11	$>$	is greater than
2.12	\geq	is greater than or equal to, is not less than
2.13	$p \Rightarrow q$	p implies q (if p then q)
2.14	$p \Leftarrow q$	p is implied by q (if q then p)
2.15	$p \Leftrightarrow q$	p implies and is implied by q (p is equivalent to q)
2.16	a	first term of an arithmetic or geometric sequence
2.17	l	last term of an arithmetic sequence
2.18	d	common difference of an arithmetic sequence
2.19	r	common ratio of a geometric sequence
2.20	S_n	sum to n terms of a sequence
2.21	S_∞	sum to infinity of a sequence

3	Operations	
3.1	$a + b$	a plus b
3.2	$a - b$	a minus b
3.3	$a \times b, ab, a.b$	a multiplied by b
3.4	$a \div b, \frac{a}{b}$	a divided by b
3.5	$\sum_{i=1}^n a_i$	$a_1 + a_2 + \dots a_n$
3.6	$\prod_{i=1}^n a_i$	$a_1 \times a_2 \times \dots a_n$
3.7	\sqrt{a}	the non-negative square root of a

3.8	$ a $	the modulus of a
3.9	$n!$	n factorial: $n! = n \times (n-1) \times \dots \times 2 \times 1$, $n \in \mathbb{N}$; $0! = 1$
3.10	$\binom{n}{r}$, nC_r , ${}_nC_r$	the binomial coefficient $\frac{n!}{r!(n-r)!}$ for $n, r \in \mathbb{Z}_0^+$, $r \leq n$ or $\frac{n!(n-1)\dots(n-r+1)}{r!}$ for $n \in \mathbb{Q}$, $r \in \mathbb{Z}_0^+$

4	Functions	
4.1	$f(x)$	the value of the function f at x
4.2	$f : x \mapsto y$	the function f maps the element x to the element y
4.3	f^{-1}	the inverse function of the function f
4.4	gf	the composite function of f and g which is defined by $gf(x) = g(f(x))$
4.5	$\lim_{x \rightarrow a} f(x)$	the limit of $f(x)$ as x tends to a
4.6	Δx , δx	an increment of x
4.7	$\frac{dy}{dx}$	the derivative of y with respect to x
4.8	$\frac{d^n y}{dx^n}$	the n th derivative of y with respect to x
4.9	$f'(x)$, $f''(x)$, ..., $f^{(n)}(x)$	the first, second, ..., n th derivatives of $f(x)$ with respect to x
4.10	\dot{x} , \ddot{x} , ...	the first, second, ... derivatives of x with respect to t
4.11	$\int y dx$	the indefinite integral of y with respect to x
4.12	$\int_a^b y dx$	the definite integral of y with respect to x between the limits $x = a$ and $x = b$

5	Exponentials and logarithmic functions	
5.1	e	base of natural logarithms
5.2	e^x , $\exp x$	exponential function of x
5.3	$\log_a x$	logarithm to the base a of x
5.4	$\ln x$, $\log_e x$	natural logarithm of x

6	Trigonometric functions	
6.1	$\left. \begin{array}{l} \sin, \cos, \tan \\ \operatorname{cosec}, \sec, \cot \end{array} \right\}$	the trigonometric functions
6.2	$\left. \begin{array}{l} \sin^{-1}, \cos^{-1}, \tan^{-1} \\ \operatorname{arc} \sin, \operatorname{arc} \cos, \operatorname{arc} \tan \end{array} \right\}$	the inverse trigonometric functions
6.3	$^\circ$	degrees
6.4	rad	radians

6	Trigonometric functions (Further Mathematics only)	
6.5	$\left. \begin{array}{l} \operatorname{cosec}^{-1}, \sec^{-1}, \cot^{-1} \\ \operatorname{arc} \operatorname{cosec}, \operatorname{arc} \sec, \operatorname{arc} \cot \end{array} \right\}$	the inverse trigonometric functions
6.6	$\left. \begin{array}{l} \sinh, \cosh, \tanh \\ \operatorname{cosech}, \operatorname{sech}, \operatorname{coth} \end{array} \right\}$	the hyperbolic functions
6.7	$\left. \begin{array}{l} \sinh^{-1}, \cosh^{-1}, \tanh^{-1} \\ \operatorname{cosech}^{-1}, \operatorname{sech}^{-1}, \operatorname{coth}^{-1} \\ \operatorname{arsinh}, \operatorname{arcosh}, \operatorname{artanh} \\ \operatorname{arcosech}, \operatorname{arcsech}, \operatorname{arcoth} \end{array} \right\}$	the inverse hyperbolic functions

7	Complex numbers (Further Mathematics only)	
7.1	i, j	square root of -1
7.2	$x + iy$	complex number with real part x and imaginary part y
7.3	$r(\cos \theta + i \sin \theta)$	modulus argument form of a complex number with modulus r and argument θ
7.4	z	a complex number, $z = x + iy = r(\cos \theta + i \sin \theta)$
7.5	$\operatorname{Re}(z)$	the real part of z , $\operatorname{Re}(z) = x$
7.6	$\operatorname{Im}(z)$	the imaginary part of z , $\operatorname{Im}(z) = y$
7.7	$ z $	the modulus of z , $ z = \sqrt{x^2 + y^2}$
7.8	$\arg(z)$	the argument of z , $\arg(z) = \theta, -\pi < \theta \leq \pi$
7.9	z^*	the complex conjugate of z , $x - iy$

8	Matrices (Further Mathematics only)	
8.1	\mathbf{M}	a matrix \mathbf{M}
8.2	$\mathbf{0}$	zero matrix
8.3	\mathbf{I}	identity matrix
8.4	\mathbf{M}^{-1}	the inverse of the matrix \mathbf{M}
8.5	\mathbf{M}^T	the transpose of the matrix \mathbf{M}
8.6	$\Delta, \det \mathbf{M}$ or $ \mathbf{M} $	the determinant of the square matrix \mathbf{M}
8.7	$\mathbf{M}\mathbf{r}$	image of column vector \mathbf{r} under the transformation associated with the matrix \mathbf{M}

9	Vectors	
9.1	$\mathbf{a}, \underline{\mathbf{a}}, \hat{\mathbf{a}}$	the vector $\mathbf{a}, \underline{\mathbf{a}}, \hat{\mathbf{a}}$; these alternatives apply throughout section 9
9.2	\overrightarrow{AB}	the vector represented in magnitude and direction by the directed line segment AB
9.3	$\hat{\mathbf{a}}$	a unit vector in the direction of \mathbf{a}

9.4	$\mathbf{i}, \mathbf{j}, \mathbf{k}$	unit vectors in the directions of the Cartesian coordinate axes
9.5	$ \mathbf{a} , a$	the magnitude of \mathbf{a}
9.6	$ \overline{AB} , AB$	the magnitude of \overline{AB}
9.7	$\begin{pmatrix} a \\ b \end{pmatrix}, a\mathbf{i} + b\mathbf{j}$	column vector and corresponding unit vector notation
9.8	\mathbf{r}	position vector
9.9	\mathbf{s}	displacement vector
9.10	\mathbf{v}	velocity vector
9.11	\mathbf{a}	acceleration vector

9	Vectors (Further Mathematics only)	
9.12	$\mathbf{a} \cdot \mathbf{b}$	the scalar product of \mathbf{a} and \mathbf{b}

10	Differential equations (Further Mathematics only)	
10.1	ω	angular speed

11	Probability and statistics	
11.1	$A, B, C, \text{ etc}$	events
11.2	$A \cup B$	union of the events A and B
11.3	$A \cap B$	intersection of the events A and B
11.4	$P(A)$	probability of the event A
11.5	A'	complement of the event A
11.6	$P(A B)$	probability of the event A conditional on the event B
11.7	$X, Y, R, \text{ etc}$	random variables
11.8	$x, y, r, \text{ etc}$	values of the random variables $X, Y, R, \text{ etc}$
11.9	x_1, x_2, \dots	values of observations

11.10	f_1, f_2, \dots	frequencies with which the observations x_1, x_2, \dots occur
11.11	$p(x), P(X = x)$	probability function of the discrete random variable X
11.12	p_1, p_2, \dots	probabilities of the values x_1, x_2, \dots of the discrete random variable X
11.13	$E(X)$	expectation of the random variable X
11.14	$\text{Var}(X)$	variance of the random variable X
11.15	\sim	has the distribution
11.16	$B(n, p)$	binomial distribution with parameters n and p , where n is the number of trials and p is the probability of success in a trial
11.17	q	$q = 1 - p$ for binomial distribution
11.18	$N(\mu, \sigma^2)$	Normal distribution with mean μ and variance σ^2
11.19	$Z \sim N(0,1)$	standard Normal distribution
11.20	ϕ	probability density function of the standardised Normal variable with distribution $N(0,1)$
11.21	Φ	corresponding cumulative distribution function
11.22	μ	population mean
11.23	σ^2	population variance
11.24	σ	population standard deviation
11.25	\bar{x}	sample mean
11.26	s^2	sample variance
11.27	s	sample standard deviation
11.28	H_0	Null hypothesis
11.29	H_1	Alternative hypothesis
11.30	r	product moment correlation coefficient for a sample
11.31	ρ	product moment correlation coefficient for a population

12	Mechanics	
12.1	kg	kilograms
12.2	m	metres
12.3	km	kilometres
12.4	m/s, m s ⁻¹	metres per second (velocity)
12.5	m/s ² , m s ⁻²	metres per second per second (acceleration)
12.6	F	force or resultant force
12.7	N	newton
12.8	Nm	newton metre (moment of force)
12.9	<i>t</i>	time
12.10	<i>s</i>	displacement
12.11	<i>u</i>	initial velocity
12.12	<i>v</i>	velocity or final velocity
12.13	<i>a</i>	acceleration
12.14	<i>g</i>	acceleration due to gravity
12.15	μ	coefficient of friction

A2

Appendix B

Mathematical formulae and identities

Students must use the following formulae and identities for AS and A-level Mathematics, without these formulae and identities being provided, either in these forms or in equivalent forms. These formulae and identities may only be provided where they are the starting point for a proof or as a result to be proved.

Pure mathematics	
Quadratic equations	$ax^2 + bx + c = 0$ has roots $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Laws of indices	$a^x a^y \equiv a^{x+y}$ $a^x \div a^y \equiv a^{x-y}$ $(a^x)^y \equiv a^{xy}$
Laws of logarithms	$x = a^n \Leftrightarrow n = \log_a x$ for $a > 0$ and $x > 0$ $\log_a x + \log_a y \equiv \log_a (xy)$ $\log_a x - \log_a y \equiv \log_a \left(\frac{x}{y}\right)$ $k \log_a x \equiv \log_a (x^k)$
Coordinate geometry	<p>A straight line, gradient m passing through (x_1, y_1) has equation $y - y_1 = m(x - x_1)$</p> <p>Straight lines with gradients m_1 and m_2 are perpendicular when $m_1 m_2 = -1$</p>
Sequences	<p>General term of an arithmetic progression: $u_n = a + (n - 1)d$</p> <p>General term of a geometric progression: $u_n = ar^{n-1}$</p>

<p>Trigonometry</p>	<p>In the triangle ABC:</p> <p>sine rule: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$</p> <p>cosine rule: $a^2 = b^2 + c^2 - 2bc \cos A$</p> <p>area: $\frac{1}{2}ab \sin C$</p> <p>$\cos^2 A + \sin^2 A \equiv 1$</p> <p>$\sec^2 A \equiv 1 + \tan^2 A$</p> <p>$\operatorname{cosec}^2 A \equiv 1 + \cot^2 A$</p> <p>$\sin 2A \equiv 2 \sin A \cos A$</p> <p>$\cos 2A \equiv \cos^2 A - \sin^2 A$</p> <p>$\tan 2A \equiv \frac{2 \tan A}{1 - \tan^2 A}$</p>
<p>Mensuration</p>	<p>Circumference (C) and area (A) of a circle, radius r and diameter d.</p> <p>$C = 2\pi r = \pi d$</p> <p>$A = \pi r^2$</p> <p>Pythagoras' Theorem: In any right-angled triangle, where a, b and c are the lengths of the sides and c is the hypotenuse:</p> <p>$c^2 = a^2 + b^2$</p> <p>Area of a trapezium: $\frac{1}{2}(a + b)h$ where a and b are the lengths of the parallel sides and h is their perpendicular separation</p> <p>Volume of a prism = area of cross section \times length</p> <p>For a circle of radius r, where an angle at the centre of θ radians subtends an arc of length s and encloses an associated sector of area A:</p> <p>$s = r \theta$</p> <p>$A = \frac{1}{2}r^2 \theta$</p>

Calculus and differential equations

Differentiation

Function	Derivative
x^n	nx^{n-1}
$\sin kx$	$k \cos kx$
$\cos kx$	$-k \sin kx$
e^{kx}	ke^{kx}
$\ln x$	$\frac{1}{x}$
$f(x) + g(x)$	$f'(x) + g'(x)$
$f(x)g(x)$	$f'(x)g(x) + f(x)g'(x)$
$f(g(x))$	$f'(g(x))g'(x)$

Integration

Function	Derivative
x^n	$\frac{1}{n+1}x^{n+1} + c, n \neq -1$
$\cos kx$	$\frac{1}{k}\sin kx + c$
$\sin kx$	$-\frac{1}{k}\cos kx + c$
e^{kx}	$\frac{1}{k}e^{kx} + c$
$\frac{1}{x}$	$\ln x + c, x \neq 0$
$f'(x) + g'(x)$	$f(x) + g(x) + c$
$f'(g(x))g'(x)$	$f(g(x)) + c$

Area under a curve $= \int_a^b y \, dx \, (y \geq 0)$

Vectors	$ \mathbf{x}\mathbf{i} + y\mathbf{j} + z\mathbf{k} = \sqrt{(x^2 + y^2 + z^2)}$
---------	---

Mechanics	
Forces and equilibrium	<p>Weight = mass \times g</p> <p>Friction: $F \leq \mu R$</p> <p>Newton's second law in the form: $F = ma$</p>
Kinematics	<p>For motion in a straight line with variable acceleration:</p> $v = \frac{dr}{dt}$ $a = \frac{dv}{dt} = \frac{d^2r}{dt^2}$ $r = \int v \, dt$ $v = \int a \, dt$

Statistics	
The mean of a set of data	$\bar{x} = \frac{\sum x}{n} = \frac{\sum fx}{\sum f}$
The standard Normal variable	$Z = \frac{X - \mu}{\sigma}$ where $X \sim N(\mu, \sigma^2)$

Get help and support

Visit our websites for information, guidance, support and resources:

aqa.org.uk

allaboutmaths.aqa.org.uk

You can also talk directly to the Maths Curriculum team

E: maths@aqa.org.uk

T: 0161 957 3852